



REVIEW ARTICLE

MicroRNAs and JAK/STAT3 signaling: A new promising therapeutic axis in blood cancers

Mehdi Sajjadi-Dokht ^a, Talar Ahmad Merza Mohamad ^b,
 Heshu Sulaiman Rahman ^{c,d}, Marwah Suliman Maashi ⁱ,
 Svetlana Danshina ^e, Navid Shomali ^{a,f}, Saeed Solali ^{a,f},
 Farough Marofi ^{a,f}, Elham Zeinalzadeh ^{a,f}, Morteza Akbari ^a,
 Ali Adili ^g, Ramin Aslaminabad ^a, Majid Farshdousti Hagh ^{a,*},
 Mostafa Jarahian ^{h,**}

^a Immunology Research Center, Tabriz University of Medical Sciences, Tabriz 5166614766, Iran

^b Department of Pharmacology, College of Pharmacy, Hawler Medical University, Erbil 44001, Iraq

^c Department of Physiology, College of Medicine, University of Sulaymaniyah, Sulaymaniyah 46001, Iraq

^d Department of Medical Laboratory Sciences, Komar University of Science and Technology, Sulaymaniyah 46002, Iraq

^e Department of Propaedeutics of Dental Diseases, I. M. Sechenov First Moscow State Medical University, Moscow 119991, Russia

^f Department of Immunology, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz 5166614766, Iran

^g Department of Oncology, Tabriz University of Medical Sciences, Tabriz 5166614766, Iran

^h German Cancer Research Center (DKFZ), Toxicology and Chemotherapy Unit (G401), Heidelberg 69120, Germany

ⁱ Stem Cells and Regenerative Medicine Unit at King Fahad Medical Research Centre, Jeddah 21589, Saudi Arabia

Received 30 April 2021; received in revised form 16 September 2021; accepted 22 October 2021

Available online 3 December 2021

KEYWORDS

Interleukin;
 JAKs;
 Leukemia;
 miRNAs;

Abstract Blood disorders include a wide spectrum of blood-associated malignancies resulting from inherited or acquired defects. The ineffectiveness of existing therapies against blood disorders arises from different reasons, one of which is drug resistance, so different types of leukemia may show different responses to treatment. Leukemia occurs for a variety of genetic and acquired reasons, leading to uncontrolled proliferation in one or more cell lines. Regarding the genetic defects, oncogene signal transducer and activator of transcription (STAT) family

* Corresponding author: German Cancer Research Center (DKFZ), Toxicology and Chemotherapy Unit (G401), Heidelberg 69120, Germany.

** Corresponding author.

E-mail addresses: m.farshdousti@gmail.com (M. Farshdousti Hagh), mostafajarahian@gmail.com (M. Jarahian).

Peer review under responsibility of Chongqing Medical University.

STAT3

transcription factor, especially STAT3, play an essential role in hematological disorders onset and progress upon mutations, dysfunction, or hyperactivity. Besides, microRNAs, as biological molecules, has been shown to play a dual role in either tumorigenesis and tumor suppression in various cancers. Besides, a strong association between STAT3 and miRNA has been reported. For example, miRNAs can regulate STAT3 via targeting its upstream mediators such as IL6, IL9, and JAKs or directly binding to the STAT3 gene. On the other hand, STAT3 can regulate miRNAs. In this review study, we aimed to determine the role of either microRNAs and STAT3 along with their effect on one another's activity and function in hematological malignancies. Copyright © 2021, Chongqing Medical University. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Abbreviations

JAK	janus kinase	TLGL	T-cell large granular lymphocytic
STAT	signal transducer and activator of transcription	ANC	absolute neutrophil count
c-myc	c-myelocytomatosis	HuR	human antigen R
Bcl-2	B-cell lymphoma 2	MEG3	maternally expressed gene 3
Mcl-1	myeloid cell leukemia 1	lncRNAs	long noncoding RNAs
Bcl-XL	B-cell lymphoma-extra large	HDAC1	histone deacetylase 1
CLL	chronic lymphocytic leukemia	DNMT	DNA methyl transferase
IL9	interleukin 9	SCA	sickle cell anemia
PBMC	peripheral blood mononuclear cell	YY1	Yin Yang 1
BCR	B-cell receptor	GATA1	GATA binding protein 1
ZAP70	zeta chain of T cell receptor-associated protein kinase 70	EPO	erythropoietin
PIAS	protein inhibitor of activated STAT3	TPO	thrombopoietin
PDCD4	programmed cell death 4	HSPCs	hematopoietic stem and progenitor CD34 ⁺ cells
PTEN	phosphatase and tensin	SOCS6	suppressor of cytokine signaling 6
MAPK	mitogen-activated protein kinase	MPN	myeloproliferative neoplasms
GFP	green fluorescent protein	AEL	acute erythroid leukemia
ALL	acute lymphoblastic leukemia	CB	cord blood
CML	chronic myeloid leukemia	LIF	leukemia inhibitory factor
		HIF1	hypoxia-inducible factor 1
		MCL	mantle cell lymphoma

Introduction

Among the intracellular defects, types of genetic mutations, deletions, unbalanced chromosomal translocation,^{1–3} and dysregulation of signal transducer factors including transcription factors are the most important underlying causes of acute leukemia disorders.^{4–7} The JAK/STAT (Janus kinase/signal transducer and activator of transcription) signaling pathway is one of the most important signaling pathways that participated in cell proliferation, differentiation, apoptosis as well as migration.^{8–11} Reports have suggested that dysregulation and overactivation of JAKs and STATs proteins is an influential marker in the development of hematological and non-hematological cancers.^{7,12–16} Therefore, suppressing or blocking this pathway is one of the significant therapeutic approaches for the treatment of leukemias in which this signaling pathway is impaired. Members of the STAT family, especially STAT3, are oncogenic factors, playing diverse roles in the biological function of cells covering proliferation, survival, angiogenesis along with metastasis, in particular, in undifferentiated cells.^{11,17–19}

The STAT family cytoplasmic transcription factor includes seven DNA-binding proteins, STAT1, STAT2, STAT3, STAT4, STAT5a, STAT5b, and STAT6 which are typically activated via phosphorylation on tyrosine residues by the JAK kinase family.²⁰ After phosphorylation, they are dimerized, and subsequently translocated into the nucleus. After that, these STAT dimers bind to specific gene promoter sequences and modify the transcription of genes involved in cellular processes adjustment, encompassing differentiation, proliferation, and apoptosis.^{8,17,21} STAT3 plays a significant role in tumorigenesis by affecting the expression of cell cycle regulators (e.g., c-Myc [cellular Myc], cyclin D1) and cancer-promoting genes such as members of the anti-apoptotic Bcl-2 (B-cell lymphoma 2) family (e.g., Mcl-1 [myeloid cell leukemia 1], Bcl-2, Bcl-XL [B-cell lymphoma-extra large]).²² To date, numerous strategies have been investigated or even used to find therapeutic strategies for suppressing JAK/STAT signaling.^{11,13,20,23,24} In continued efforts to more understanding the mechanisms of JAK/STAT3 signaling inhibition, a large number of studies have identified the pivotal role of miRNAs in the regulation of STAT3 or signaling

pathway in which STAT3 is involved.^{25–29} On the other hand, several studies have implied the possible role of STAT3 in regulating miRNA expression.^{30–32}

miRNAs are an endogenous and non-coding class of single-stranded RNAs containing 19 to 25 nucleotides, modifying gene expression post-transcriptionally through binding to the 3' untranslated regions (3' UTRs) leading to the promotion of the mRNA degradation or triggering the inhibition of mRNA translation.^{33–35} miRNAs actively contribute to a variety of cellular processes such as the production of auto-antibodies, cellular immunity, cell proliferation, differentiation, cell cycle, and apoptosis in concomitant with association with various other events such as neurodevelopment, DNA editing-repairing, and oxidative stress.^{36–40} It has been also robustly evidenced that the dysregulation of miRNAs expression is involved in the tumor microenvironment (TME) and drug resistance.^{41–43} In Figure 1, microRNA biogenesis has been shown.

In sum, numerous studies have shown that miRNAs and STAT3 may stimulate a fundamental regulatory effect on each other directly or indirectly.^{44–47} Herein, we have highlighted the potent role of miRNAs in the regulation of STAT3 expression and function in hematological disorders.

Interaction between miRNAs and JAK/STAT3 signaling in chronic lymphocytic leukemia (CLL)

Chronic lymphocytic leukemia (CLL) is one of the most common hematological disorders in the world and the most common adult leukemia disorder in the western hemisphere.⁴⁸ Generally, about 95% of cases are B cells⁴⁹ typically characterized by the accumulation of mature and small B cell phenotype expressing CD5⁺ and CD19⁺ immune phenotype markers, affecting peripheral blood, bone marrow (BM), and lymphatic tissues.⁴⁸

Relationship between IL9, STAT3, miR-A21, miR-155 in CLL

Previous studies have displayed that IL9 is commonly increased in CLL patients.⁴⁸ Chen et al also observed that the proliferation of CLL cells was increased after using recombinant human IL9, while apoptosis was decreased. They also analyzed the relationship between the upregulation of IL9 expression and the expression levels of STAT3, P-STAT3, miR-21, miR-51 in peripheral blood mononuclear cell (PBMC) of CLL patients, and found an increase in expression of STAT3 and P-STAT3 in CLL patients.⁴⁸ IL9 is more commonly known as Th2 cytokine, contributes to allergic diseases.⁵⁰ Findings have presented that IL9 in addition to the involvement in T-reg and mast cell-mediated tumor immunity,⁵¹ may participate in growth, tumor progression, and the anti-apoptotic process.⁴⁸ Many studies have shown that the IL9- α chain stimulates mutant JAK1 phosphorylation resulted in activation of the STAT family, in particular, STAT3.^{52–54} Chen and his coworkers signified that the transfection of miR-21 and miR-21 overexpressed CLL cell lines could stimulate IL9 production in these cells, enabling IL9 production in CLL cells upon promotion of p-STAT3

cellular levels.⁴⁸ These findings implied that an IL9 endogenous/IL9 exogenous/miR-21/miR-155/STAT3 axis exists in CLL cells, which could be useful in finding new therapeutic patterns.⁴⁸

The relationship between miR-21 and the STAT3 signaling pathway in CLL cells

CLL cells are highly dependent on their microenvironment and the B-cell receptor (BCR) signaling pathway plays an important role in this regard.^{49,55} Among the proteins involved in cell-to-microenvironment interaction, the zeta chain of T cell receptor-associated protein kinase 70 (ZAP70) can improve cell response to microenvironment stimuli.⁴⁹

miR-21 is overexpressed in various leukemic disorders including CLL,^{56,57} possibly involved in the development of drug resistance and survival along with augmenting of disease progression.⁵⁸ Besides, the presence of the relationship between miR-21 and poor prognosis in CLL⁵⁹ as well as cell proliferation and oncogenesis has been evidenced.⁶⁰ Carabia et al showed that stimulation of BCR signaling by the microenvironment can regulate the expression of miR-21 and its target repressor genes, including protein inhibitor of activated STAT3 (PIAS3),⁶¹ programmed cell death 4 (PDCD4),⁶² and phosphatase and tensin homolog (PTEN)⁶³ via the signaling pathway stimulated or progressed by mitogen-activated protein kinase (MAPK or MAP kinase) and STAT3.⁴⁹ The ZAP70 protein plays an important role in different signaling pathways also modulates the interaction between the cells and related microenvironments.⁴⁹ Carabia and her colleagues analyzed the expression changes of miR-21 following ZAP70 status in CD19⁺ B cells derived from patients who were diagnosed with CLL and found that miR-21 was highly expressed in patients with higher expression of ZAP70. Also, the study of miR-21 expression level in Ramos cell (human Burkitt's lymphoma B cells) transfected with GFP-ZAP70 demonstrated similar results.⁴⁹ In another study, the stimulation of the IL-6 receptor (one of the B cell receptors) on the surface of the myeloma B cells resulted in pre-miR regulation by STAT3 translocation into the nucleus.⁶⁴ Accordingly, Carabia et al identified that the stimulation of the BCR signaling pathway caused ZAP-70 activation, which in turn, triggered an enhancement in mature miR-21 levels via inducing the STAT3 binding to the pre-miR-21 transcript promoter.⁴⁹

miR-155 and its association with STAT3 signaling in CLL cells

The miR-155 is involved in tumorigenesis and autoimmunity⁶⁵ and its overexpression can lead to lymphoma onset in mice.⁶⁶ The miR-155 regulates the proliferation and development of hematopoietic cells²¹ as well as contributed to immune cell response, production of antibodies, cytokines, and antigen expression.^{67,68} In this context, a study showed that overexpression of miR-155 in the murine model leads to B cell proliferation and is associated with lymphoma development.⁶⁶ Similarly, overexpression of miR-155 has been observed in CLL^{56,69,70} accompanied by




Figure 1 miRNA biogenesis. The majority of genes, which encode the miRNAs, are in promoter regions of intronic genes. Firstly, RNA polymerase II transcribes the miRNA genes to the primary miRNA (pri-miRNA) form. Drosha, a type III RNase, incorporated with the cofactor protein DGCR8, binds to the primary miRNA (pri-miRNA) transcript and cleaves it, in the nucleus to form precursors-miRNAs. The pre-miRNAs are exported to the cytosol by Exportin 5 (whereas in the drosha independent pathway pri-miR are exported into the cytosol by Exportin 1 and then cleaved to pre-miRNAs form), and subsequently cleaved by Dicer (a type III RNase) to miRNA duplex form incorporation with TAR RNA binding protein (TRBP). The argonaute (AGO) family and its associated protein mediate the processing of the miRNA duplex. After unwinding and strand selection, the mature miRNA can target recognition. Next, mature miRNA binds to the RISC and so it is capable of targeting the mRNAs via complementary sites and results in translational repression or mRNA degradation.

Hodgkin and non-Hodgkin's lymphoma.^{68,71,72} Other investigations have implied that IL-6 may activate STAT3 expression and function in CLL cells^{73,74}; on the other hand, overexpression of miR-155 has been found in CLL.^{56,69,70,75,76} Considering the results, it seems that IL6 increases the miR-155 expression, delivering proof of the concept of the important influences of miRNA in leukemia

pathogenesis or therapy.²¹ In this regard, Li et al detected miR-155 overexpression in CLL cells upon exposure with rh IL-6 and verified STAT3 binding to the miR-155 promoter in rh IL-6-exposed CLL cells. Despite the direct regulation of miR-155 transcription by STAT3, they suggested that IL-6 increased STAT3 binding to the miR-155 promoter. Structural and molecular analysis verified the existence of the

two binding sites in the miR-155 gene promoter for STAT3.²¹ STAT3, after phosphorylation, joins to the gamma interferon activation site (GAS)-LIKE components located in the promoter region of various genes.⁷⁷ Indeed, Li et al introduced two GAS-LIKE elements inside the miR-155 promoter, enabling STAT3 binding to the miR-155 promoter by these elements.²¹ Thus, they confirmed that IL-6 induces miR-155 activation in CLL cells via STAT3, and consequently regulates hematopoietic cell proliferation.²¹

Interaction between miRNAs and JAK/STAT3 signaling in acute lymphoblastic leukemia (ALL)

Acute lymphoblastic leukemia (ALL) is a heterogeneous hematologic disease characterized by the high proliferation of non-mature lymphocytic cells in BM, peripheral blood, and other organs. About 4000 people have been involved annually in the United States that 75%–80% of them are children.⁷⁸

Relationship between miR-451a and JAK/STAT3 signaling in Philadelphia-positive ALL (Ph⁺ ALL)

ALL with BCR-ABL1 gene fusion (Philadelphia chromosome-positive) in the precursor of B-lineage subtypes, which is resulted in constitutive activation of ABL tyrosine kinase, is one of the most fatal leukemia showing unfavorable prognosis^{79,80}; however, commonly demonstrate an appropriate response to tyrosine kinase inhibitors (TKI).^{79,80} Nevertheless, the relapses incidence rate is notable in these patients, highlighting the importance of new insights to treating this condition.⁸¹ It has been clarified that BCR/ABL fusion protein in chronic myeloid leukemia (CML), stimulates the expression of IL-6, which is a multipotent cytokine that prepares the favorable micro-environment for expansion and maintenance of the leukemic stem cells.⁸² Despite that BCR/ABL fusion gene is the indicator marker of Ph⁺ ALL, the relation between BCR/ABL fusion and IL-6 in Ph⁺ ALL has not yet been elucidated entirely.⁸³ As described, IL-6 binds to its specific receptor (IL-6R) and then triggers the JAK/STAT3 signaling pathway.^{84,85} In a study, Jiang et al evaluated the level of BCR-ABL expression in Ph⁺ ALL patients to investigate the association between BCR-ABL and miR-451 in mononuclear cells of these patients, and found that BCR-ABL has an inverse relation with miR-451.⁸³ Bioinformatic analysis showed that IL-6R can be a target of miR-451a.⁸⁶ It has been reported that miR-451a represses the proliferation of some types of cancer cells such as lung adenocarcinoma⁸⁷ and renal cell carcinoma.⁸⁸ Besides, in CML, miR-451a is inversely related to BCR-ABL gene expression levels, and its downregulation is paralleled to imatinib resistance.⁸⁹ Examination of the underlying mechanisms of miR-451a on the ALL cell line with Philadelphia chromosome-positive showed that this mi-RNA directly suppresses IL-6R via targeting its 3'-UTR. Jiang et al reported that BCR/ABL mRNA levels were inversely correlated to miR-451a and positively associated with serum IL-6 concentration.⁸³ They also indicated that miR-

451a downregulated the levels of phosphorylated JAK2 and STAT3 in the absence of any impact on total rates of JAK2 and STAT3, indicating the inhibitory effects of the miR-451 on IL6/JAK/STAT3 axis upon suppressing the IL6R and p-JAK and p-STAT3 downstream molecules.⁸³ Thereby, it has been suggested that miR-451 is downregulated by BCR-ABL at Ph⁺ ALL patients to provide a survival advantage in leukemic cells via the IL6/JAK/STAT3 pathway. Therefore, elevation in miR-451 expression in these patients could rise the promise for the treatment of this acute and severe type of leukemia.⁸³

The relationship between JAK/STAT pathway and miR-146b and FASL for inducing neutropenia in T-cell large granular lymphocytic (TLGL) leukemia

T-cell large granular lymphocytic (TLGL) is a rare type of leukemia characterized by the abnormal proliferation of large granular T lymphocytes in peripheral blood.^{90,91} TLGL leukemic cells are divided into two clusters, including cluster A that showed CD8⁺/CD4[−]/CD57⁺ immune phenotype, and cluster B that showed CD4⁺/CD8[−]/CD57⁺ immune phenotype.⁹² The underlying cause of TLGL is not fully well known, an impairment in apoptotic signals and increase in survival involve be considered as possible pathogenesis of these types of blood malignancies.^{92,93} Among these, JAK/STAT signaling is one of the well-known involved pathways in TLGL progress especially in the CD8⁺/CD4[−] immunophenotype which is associated with neutropenia.^{92,94,95}

The mechanism of neutropenia in TLGL is not fully understood, while degeneration of mature neutrophils and myeloid progenitors through FAS/FASL signaling has been suggested as a potent corresponding mechanism.⁹² Previously, it has been reported that promoted levels of soluble FASL in the blood circulation are one of the most important possible factors contributing to the pathogenesis of neutropenia in T-LGL patients.^{96,97} Many studies also have the existence of the improved concentrations of soluble FASL in TLGL,^{95,96,98,99} which supports neutropenia occurrence in TLGL patients. It has been observed that about 35% of TLGL patients have a hotspot mutation in STAT3, supporting the overactivation of STAT3.^{100,101} Teramo et al have reported that STAT3 plays a key role in FASL transcription and showed that a promoted level of STAT3 activity is associated with a higher rate of FASL expression in TLGL patients⁹⁵; however, the molecular mechanism of FASL regulation by STAT3 has not yet been elucidated. In another study, Mariotti and her co-workers also detected elevated levels of tyrosine-phosphorylated (YP)-STAT3 in CD8⁺ T-LGL patients, but not in CD4⁺ T-LGL patients.⁹² Mariotti et al revealed that FASL mRNA expression is correlated with the upregulation of STAT3 activation and inversely with the absolute neutrophil counts (ANC).⁹² They determined that miR-146b was downregulated in CD8⁺ TLGL compared to CD4⁺,⁹² as well as found that there is an association between miR-146b expression and ANC levels concomitant with the levels of YP-STAT3 in T-LGL patients.⁹² In addition, they found that miR-146b has an inverse correlation with STAT3 tyrosine phosphorylation, neutropenia, FASL

expression, and soluble FASL release in blood circulation,^{92,95} emphasizing the importance of the STAT3-miR146b-FasL axis in TLGL leukemia.⁹² Respecting the previous findings that STAT3 can stimulate inhibitory influence on gene expression through inducing the target genes promoter methylation,^{102,103} it has been confirmed that STAT3 stimulates miR-146b promoter methylation through regulating the expression of methyl transferase-1 in solid tumors and T lymphocytic malignancies.¹⁰⁴ Regardless of the presence of correlation between STAT3 function and miR-146b cellular levels in malignant condition,^{104,105} it also has been reported STAT3 activates miR-146b in normal tissues.^{106,107} Overall, Mariotti et al suggested that STAT3 suppressed the miR-146b expression in TLGL by inducing the miR-146b promoter methylation.⁹² Considering molecular analysis, human antigen R (HuR), which plays a well-known role in mRNA stabilization and FASL expression, has been shown that can be another target of miR-146b.^{108–110} Mariotti et al detected that HuR protein is an endogenous target of miR-146b in CD8⁺ T-LGLs, and also indicated that miR-146b downregulated the FASL expression indirectly and post-transcriptionally through reducing the HuR protein levels. Accordingly, they suggested that consistent over-activity of STAT3 in CD8⁺ T-LGLs resulted in the loss of miR-146b, which in turn led to the HuR protein translation, enabling FasL production and neutropenia incidence.^{92,95}

Interaction between miRNAs and JAK/STAT3 signaling in CML

CML is a hematopoietic stem cell clonal disorder characterized by the generation of the Philadelphia chromosome (Ph⁺) due to the BCR-ABL oncogenic protein fusion resulting from translocation t(9;22) and comprises about 15% of leukemia diagnosed in adults.¹¹¹

The relationship between miR-147 and STAT3 signaling

The miR-147 has been identified about various cancers^{112–115} through showing dual roles, including tumorigenesis and tumor suppressor roles, in a variety of human cancers. For example, upregulation of miR-147 has been observed in human gastric and liver cancers, whereas its downregulation has been reported in human breast cancer.¹¹⁶ In a study, Han et al showed that the hypoxia-induced damage was intensified in the PC12 cell line, commonly derived from a transplantable rat pheochromocytoma, due to the attenuation in miR-147 levels.¹¹⁷ In detail, they showed that maternally expressed gene 3 (MEG3) as a long noncoding RNAs (lncRNAs) boosted hypoxia injury in PC12 cells through down-regulating miR-147, as well as demonstrated that the overexpression of MEG3 and miR-147 induced apoptosis and inhibited target cells proliferation.¹¹⁷ lncRNAs and miRNAs usually act as competing endogenous RNAs. Evaluation of the miRNAs database to identify MEG3 interacting miRNAs has revealed that miR-147 is one of the potential miRNAs in this regard. Investigations revealed that the levels of MEG3 and miR-147 expression in the KCL22 and K562, two human CML cells,

were decreased in comparison to healthy BMMCs. It seems that MEG3 can bind to miR-147 and eventually reduce miR-147 expression. Li et al also observed a high level of methylation of miR-147 and MEG3 in CML patients but not in healthy individuals. Accordingly, they showed that the expression rate of the methylation-related genes, such as HDAC1 (histone deacetylase 1), DNMT1 (DNA methyl transferase 1), DNMT3A, DNMT3B, were remarkably promoted in the CML patients with accelerated phase. Therefore, they concluded that the downregulation of miR-147 and MEG3 levels in the CML patients was probably mediated by histone deacetylation and DNA methylation. Besides, they suggested that JAK2 and STAT3 can negatively regulate MEG3 upon binding to it, as well as described that MEG3 can downregulate STAT3 by suppressing the phosphorylation of JAK/STAT.¹¹⁸ In sum, they suggested that there is a negative feedback between MEG3 and STAT3. On the other hand, they showed that miR-147 and MEG3 can regulate each other negatively and adjust leukemia development, verifying that STAT3 and miR-147 indirectly affect each other through MEG3.¹¹⁸

The relationship between miR-574-3P and JAK/STAT3 signaling

The significant role of miR-574-3p in some cancers has been reported. For instance, miR-574-3p commonly is decreased in bladder cancer cells whereas overexpression of miR-574-3p remarkably prohibited bladder cancer cell proliferation, invasion, and migration.¹¹⁹ In addition to the miR-574-3p importance as a potential prognostic marker for breast cancer,¹²⁰ its lower expression is noticed in the early stages of gastric cancer; however, upregulation of miR-574-3p inhibits proliferation, invasion, as well as the migration of human gastric cancer cells, line.¹²¹ It is also reported that miR-574-3P has a tumor-inducing role in human osteosarcoma,¹²² while triggers an inhibitory impact on esophageal squamous cell carcinoma,¹²³ implying its potentially reciprocal role in a variety of human cancers.

Yang et al showed that the miR-574-3P expression was significantly decreased in the peripheral blood of CML patients compared to the healthy donor. They also showed that overexpression of miR-574-3P resulted in significant inhibition of cell proliferation of human K562 CML cells and induced apoptosis, whereas suppression of miR-574-3P had a contrasting effect. They checked the TargetScan-Human database and found that the IL6 can be a potential direct target for miR-574-3p. They also found that the IL6 at mRNA and protein levels was remarkably inhibited upon miR-574-3p transfection. These data showed that IL6 is a direct target for miR-574-3p and IL6 expression was negatively regulated by miR-574-3p. As known, IL6 is identified as an inflammatory cytokine involved in the pathogenesis of hematological disorders such as multiple myeloma.¹²⁴ Maeda et al showed that IL6 may contribute to both myeloid proliferation and lymphocytopenia.¹²⁵ They suggested that IL6 could control the cell destination of leukemic multipotent progenitor cells and may support a positive feedback loop to maintain CML progression.⁸² It is also known to be a prognostic factor for the follow-up of imatinib treatment in CML patients.¹²⁶ It has already been

found that overexpression of IL6 significantly induces K562 CML cell proliferation, and conversely inhibits their apoptosis.¹²⁷ Also, reports confirm that IL-6 can activate the JAK/STAT3 and MAPK signaling pathway and is involved in the development of CML.⁵⁵ Rendering findings, Yang et al proposed that miR-574-3P can suppress the IL6/JAK/STAT3 signaling pathway through directly targeting IL6, which in turn, inhibits the proliferation and induce apoptosis in CML cells,¹²⁷ describing miR-574-3p as a potent target for CML treatment.

The relationship between miR-34a and embryonic hemoglobin production in myeloid cells mediated by STAT3

Sickle cell anemia (SCA), which is characterized by abnormal hemoglobin S (Hb S) production, is one of the most common genetic hematological disorders. Annually more than 300,000 newborns are born with cyclic anemia, which is one of the most common blood disorders in the world associated with a poor prognosis.¹²⁸ Despite the recent increase in new therapeutic approaches, hydroxyurea is still the principal of SCA treatment showing the competence to reduce SCA-related mortalities.¹²⁹ Indeed, the most effective treatment for the cycle cell anemia is increasing Hb F production with the formula $\alpha_2\gamma_2$ (a combination of two alpha chains and two gamma chains globulin) that improves clinical symptoms and consequently prolongs the patient overall survival rates.^{130,131} The therapeutic role of Hb F is related to its inhibitory effects on Hb S polymerization,¹³² which is one of the main pathogens and risk factors in patients with SCA.¹³³ Importantly, comprehensive studies have indicated that a wide spectrum of genes is targeted by miR-34a, including the genes known as a repressor of gamma-globin, a variant of globulin chain used in hemoglobin F production, such as Yin Yang 1 (YY1),

HDAC1, and STAT3.^{75,134–137} Scientific related software predicted the binding site for miR-34a at the 3' UTR of STAT3,¹³¹ and also studies have shown that miR-34a interacts with STAT3 in K562 CML cells.¹³⁷ Interestingly, reports have revealed that GATA1 (GATA binding protein 1) and STAT3 compete for binding to the 5' UTR of the gamma globulin gene and an increase in GATA1 expression reverses gamma globulin gene silencing induced by STAT3.¹³⁸ Besides, it has been shown that during the erythroid differentiation progresses, erythropoietin (EPO) can activate STAT3 through phosphorylation,¹³⁹ leading to the inhibition of gamma globulin gene expression.¹³⁸ Therefore, these data verify that STAT3 has a function in gamma globulin gene regulation. In a study, Ward et al conducted a study to determine whether miR-34a can regulate gamma-globin gene expression through targeting the negative gamma globulin regulator genes including STAT3. Achieved consequences exhibited that the total and phosphorylated STAT3 levels were significantly decreased in miR-34a-transfected K562 cells.¹³¹ They also observed that the gamma-globin mRNA and Hb F protein levels were augmented in response to overexpression of miR-34a. Respecting that the overexpression of miR-34a leads to a reduction in STAT3 and an increase in gamma-globin and Hb F expression, WARD et al reports confirmed that likely exist an indirect mechanism for gamma-globin regulation by miR-34a via STAT3 gene silencing.¹³¹ They showed that miR-34a induced the production of Hb F in K562 cells by reducing total STAT3 and phosphorylated STAT3 levels, which have a role in silencing the gamma globulin gene.¹³¹

Interaction between miRNAs and JAK/STAT3 signaling in AML

AML is the most common acute leukemia in adults, with about 20,000 cases per year in the United States alone.¹⁴⁰




Figure 2 Role of activated STAT3 in tumor promotion. STAT3 is activated by binding to DNA and by increasing the expression of various genes involved in various biological processes such as proliferation, tumor progression, metastasis, etc., causing tumor growth.

Association with miR-494-3p overexpression and promoting the megakaryocytopoiesis in primary myelofibrosis hematopoietic stem/progenitor cells

Thrombopoietin (TPO) is produced in the liver and acts as the main regulator in megakaryopoiesis and participates in megakaryocyte proliferation and differentiation. TPO has been identified as a pan-hematopoietic cytokine and is crucial for the preservation and survival of hematopoietic stem cells.¹⁴¹ It binds to its receptor, TPOR, on the surface of hematopoietic stem and progenitor CD34⁺ cells (HSPCs) and results in the activation of JAK2, which leads to STAT3 phosphorylation and induction of megakaryopoiesis.¹⁴² Studies have supported overexpression of miR-494-3P in HSPCs in patients with primary myelofibrosis (PMF).¹⁴³ Rontauroli et al in their study showed that an increase in expression of miR-494-3P, enhanced hematopoiesis in normal hematopoietic cells.¹⁴⁴ Investigations have recommended that the suppressor of cytokine signaling 6 (SOCS6) is a target for miR-494-3p. Also, a previous study reported that SOCS6 is downregulated in HSPCs derived from PMF patients, which concurrently showed miR-494-3p upregulation.¹⁴³ SOCS6 is an important factor in the negative regulation of the JAK/STAT signaling pathway,¹⁴⁵ as well as

contributes to the mechanism of myeloproliferative neoplasms (MPN) pathogenesis.¹⁴⁴ Other investigations have evidenced that SOCS6 decreases the nuclear level of the STAT3 and attenuates its phosphorylation, supporting STAT3 downregulation.^{145,146} Rontauroli et al reported that the expression level of MK antigen, including CD41 or later CD42b antigen, has superiority in CD34⁺ cells transfected with miR-494-3p over normal cells. These results confirmed that miR-494-3p is involved in megakaryocyte differentiation in the pathogenesis of PMF. Furthermore, STAT3 may affect TPO signaling and megakaryocytopoiesis in HSPCs.¹⁴² The results of the study of Rontauroli et al demonstrated that transfection of K562 and CB (cord blood-derived) CD34⁺ cells with miR-494-3P decreased SOCS6 protein levels and subsequently increased STAT3 phosphorylation, ultimately leading to megakaryocyte hyperplasia observed in PMF patients.¹⁴⁴

Cluster miR-23a, miR-27a, miR-24 and association with JAK/STAT signaling pathway in acute erythroid leukemia (AEL)

Acute erythroid leukemia (AEL) (termed AML M6), is a rare subtype of AML with accounts for about 5% of all AML cases.^{147,148} The survival and prognosis of AEL patients are




Figure 3 Stimuli that affect the STAT3 signaling pathway. STAT3 signaling cascade can be activated by various stimulators such as cytokines and growth factors.




Figure 4 The signals mentioned in the study in which STAT3 and miRNAs are involved.

too worse than in other AML subtypes.^{149,150} As described, STAT3 plays an important role in the progression of erythroleukemia through suppression of erythroid differentiation.¹⁵¹ In a study, Su et al showed that miR-23a, miR-27a, miR-24 synergistically targeted the members of the GP130/JAK1/STAT3 signaling pathway. They also showed that miR-23a, miR-27a, and miR-24 clusters were downregulated in AEL patients. They assumed that miR-23a, miR-27a, miR-

24 cluster jointly targeted the GP130/JAK1/STAT3 pathway in AEL cells and induced the differentiation of these cells.¹⁵² Other studies have presented that the JAK1 binding to the GP130 transmembrane protein and succeeding activation of STAT3 enables signaling network formation among GP130, JAK1, and STAT3.⁷ They concluded that overexpression of miR-23a, -27, and -24 can trigger apoptosis and inhibit the deregulated proliferation of AEL cells through the

Table 1 Some of the cancers in which miRNAs and STAT3 regulates each other directly or indirectly.

Cancers	miRNA	Interaction between miR-503-5p and STAT3	Suggested pathway or signaling involved	References
Metastatic or paclitaxel-resistant ovarian cancer cells	miR-503-5p	miR-503-5p binds directly to the 3' UTRs of CD97	miR-503-5p inhibits the CD97-mediated JAK2/STAT3 pathway	190
Early gastric cancer	miRNA-200 Family	miR-200 family members were upregulated by IL11/STAT3 signaling	IL11/STAT3-dependent manner	191
Ovarian cancer	miRNA-92	STAT3 directly regulates miR-92a	STAT3/miRNA-92/DKK1/Wnt/b-catenin signaling pathway	192
Colorectal cancer	miR-34a	IL6R is a direct target of miR-34/STAT3 and JAK 2 can be potentially miR-34a targets	IL-6R/STAT3/miR-34a feedback loop	193
Human gastric cancer	MicroRNA-143	miR-143 directly targets STAT-3	Proliferation, migration, and invasion signaling in which STAT3 is involved	
Cervical cancer	miR-411	miR-411 inhibited cervical cancer progression by directly targeting STAT3	Proliferation and invasion signaling in which STAT3 is involved	26
Pancreatic cancer	miR-301a	Targeting SOCS5	miR-301a induces pancreatic cancer invasion and metastasis via JAK/STAT3 signaling by targeting SOCS5	194
Breast cancer	MicroRNA-204	Targeting JAK2	MicroRNA-204 targets JAK2 and induces cell apoptosis through the STAT3/BCL-2/survivin pathway	195
Human renal carcinoma cells	miRNA-133b/ miR-135a	miR-133b and miR-135a may target directly the 3'-UTR of JAK2	miR-133b and miR-135a induce apoptosis signaling cascade involving JAK2, STAT3, and Bcl-2	196
Pancreatic cancer cells	miRNA let-7	let-7 upregulates SOCS3 expression in PDAC cells	miRNA let-7/SOCS3/STAT3	197
Natural killer T cell cytotoxic activity in cervical cancer	miR-124-3p	Down-regulation of miR-124-3p promotes cervical cancer progression through targeting STAT3	LINC00240/microRNA-124-3p/STAT3/MICA axis	198
Epigenetic switch between inflammation and colorectal cancer	miR-181b	STAT3 up-regulates miR-181b through directly binding	STAT3-miR-181b- CYLD- NF- κ B	199
Breast cancer cells	miR-146b	miR-146b was reported as a direct target of STAT3	NF- κ B/IL-6/STAT3 pathway	104
AOM/DSS mice model induced inflammatory bowel disease	miR-221	miR-221 directly targets the PDLIM2 gene that is known to be essential for restraint NF- κ B	miR-221/PDLIM2 gene/NF- κ B/STAT3	200
Colitis-associated cancer	miR-155	miR-155 directly targets and inhibits the SOCS1(the inhibitor of STAT3)	miR-155/NF- κ B-STAT3	200
Lung cancer cells	miR-206	miR-206 decreased the angiogenesis by targeting 14-3-3z	Locking the 14-3-3z/STAT3/HIF-1a/VEGF signaling	201
Colorectal cancer cells	miR-375	JAK2, MAP3K8, and ATG7 are predicted target genes for miR-375	JAK2/STAT3 and MAP3K8/ERK signaling pathways	202
Gastric cancer cells	MicroRNA-216a	miR-216a directly targets JAK2	JAK2/STAT3 pathway	203
Papillary thyroid cancer	miRNA-148a	microRNA-148a promotes	STAT3 and PI3K/AKT signaling	204

Table 1 (continued)

Cancers	miRNA	Interaction between miR-503-5p and STAT3	Suggested pathway or signaling involved	References
Natural killer T cell cytotoxic activity in cervical cancer	microRNA-124-3p	cancer cell growth by targeting PI3K signaling pathway miR-124-3p is a direct target of LINC00240/Inhibition of miR-124-3p promotes cervical cancer progression via targeting STAT3	pathways LINC00240/microRNA-124-3p/STAT3/MICA axis	198
Breast cancer cell lines	miR-146b	a gene encoding the miR-146b is a direct STAT3 target gene/STAT3 induces miR-146b, a negative regulator in cancer/miR-146b links STAT3 to NF-kB and IL-6 as part of a negative feedback loop	miR-146b/NF-kB/I L-6/STAT3 Axis	104
Lung cancer	MicroRNA-218	miR-218 negatively regulates IL-6 receptor and JAK3 gene expression by directly targeting the 3'-UTR of their mRNAs	IL-6/STAT3 signaling pathway	205

suppressing of the GP130-JAK1-STAT3 cascade. They also reported that these miRNAs cooperatively induce erythroid differentiation via inhibiting the GP130 and suppression of the JAK1-STAT3 phosphorylation in human leukemic HEL and K562 cell lines as well as cord blood (CB)- CD34⁺ HSCs.¹⁵²

Relationship between miR-155 and JAK/STAT3 signaling and LIFRa

Laboratory studies on leukemic cells, including HL60, show that STAT3 overexpression and leukemogenesis is depending on the phosphorylation of STAT3.¹⁵³ While inhibition of STAT3 signaling leads to the induction of apoptosis in leukemic cells.¹⁵⁴ The leukemia inhibitory factor (LIF), as known as one of the members of IL-6 family cytokines, has been reported that can induce differentiation of the M1 murine myeloid leukemic cell line. It is named LIF due to its ability to induce differentiation M1 myeloid leukemic cells.¹⁵⁵ This protein executes its biological activity through its receptor located on the cell surface and a membrane-associated transducer (termed LIFR- α).¹⁵⁶ Former studies determined that STAT3 can be activated by IL-6 family cytokines,¹⁵⁷ and also revealed that IL6 family/STAT3 signaling may affect differentiation of stem and leukemic cells and other cells.^{158–161} LIFR α has been determined that bind to the GP-130 on the surface of leukemic cells and form heterodimers, which are capable of STAT3 activation.¹⁶² Similar to LIFRa, the fusion protein containing the cytoplasmic functional domain of LIFRa, like CT3 in TAT-CT3 fusion, a peptide domain vector, can induce STAT3 activation in HL60 cells.¹⁶³ The LIFRa-CT3 fusion transfection in HL60 cells resulted in suppression of the

proliferation and promoting the differentiation in HL60 myeloid cells *in vitro*.¹⁶³ Other studies have proposed that miR-155 is a primary transcript of the third exon of the B cell integration cluster (BIC) gene¹⁶⁴ typically overexpressed in the BM of patients with special subtypes of AML.^{165–167} Other examinations have represented that IL-10 suppresses miR-155 through STAT3 activation.¹⁶⁸ Moreover, XU et al reported that the TAT-CT3 fusion protein inhibits miR-155 expression following targeting STAT3 in HL-60 cells.¹⁵⁵ They determined that the TAT-CT3 fusion protein negatively regulated miR-155 expression, which is overexpressed in some type of AML, through STAT3 direct binding to miRNA gene promoter.¹⁵⁵ Also, miR-155 negatively regulates SOCS-1 in AML cells, known as the main negative regulator for the JAK/STAT signaling pathway.¹⁶⁹ XU and his colleagues showed that the TAT-CT3 fusion protein can induce differentiation in HL60 myeloid cells. They showed that TAT-CT3 transfection downregulated the miR-155 expression, which in turn, increased SOCS-1 and subsequently decreased STAT3 phosphorylation results in differentiation of leukemia cells.¹⁵⁵

The miR-17 and miR-20a and association with STAT3 and hypoxia-inducible factor 1 (HIF1)

Hypoxia-inducible factor 1 (HIF1) is an important transcription factor in response to hypoxic conditions containing two subunits including, an alpha subunit (HIF-1 α , oxygen-sensitive subunit) and a beta subunit (HIF-1 β).^{170,171} It also has a role in cancer biologics such as tumor growth, metastasis, and angiogenesis.¹⁷² Consistent with the previous studies,^{173,174} He et al showed that hypoxia

Table 2 The hematological malignancies that is affected by miRNA in relation with STAT3.

miRNA	Direct target of miRNA	MiRNA expression status in malignancy	Effect on STAT3	Type of hematological disorder	Samples	Function	Suggested pathway or signaling involved	Ref
miR-21/miR-155	IL9	Up regulated	N/A	CLL	MEC1 CLL cell line	pSTAT3 up regulated miR-155 and miR-21	Extra cellular IL9/ pSTAT3/miR-155/ miR-21/intra cellular IL9' positive feedback system in CLL cells	48
miR-21	repressor genes, such as PIAS3 and PDCD4 and PTEN	Up regulated	Indirectly via STAT3 suppressor gene	CLL	RAMOS B cell/ UE6E7T2/bone marrow stromal cell	STAT3 up regulated miRNA-21	BCR/ZAP70/STAT3/ miR-21/PTEN, PDCD4 and PIAS3	49
miR-155	N/A	Up regulated	N/A	CLL	Human multiple myeloma MM1 cells	STAT3 up regulated miR-155	IL6/STAT3/miR-155	21
miR-451a	IL6R	Down regulated	Direct binding to down-stream of IL6R as a STAT3 activator	Ph ⁺ ALL	BM-MNCs/Ph ⁺ ALL cell line SUP-B15/Ph- ALL cell line Nalm6	miR-451a downregulated pSTAT3 indirectly	miR-451a/IL6/JAK/ STAT3 pathway	83
miRNA-146b	HuR protein	Down regulated	N/A	CD8 TLGLs	PBMCs from TLGLs patient/Jurkat cells	STAT3 downregulated miRNA-146b expression	STAT3/miR146b HuR/ FASL axis	206
miRNA-147	N/A	Down regulated	miR-147 indirectly affects on STAT3 via affecting on MEG3	CML	KCL22/K562 CML cells/BMMCs	miR-147 can probably regulate STAT3 via MEG3	negative feedback loop between MEG3 and STAT3	118
MiRNA-574-3P	IL6	Down regulated	Indirectly via Increase in IL6	CML	K562 CML cells	miRNA-574-3P down regulated JAK/STAT3 signaling via IL6	miR-574-3p/IL6/JAK/ STAT3	127
miRNA-34a	STAT3	—	Directly increase in STAT3	Sickle cell anemia	K562 CML cells	miRNA-34a down regulated STAT3	miRNA-34a/STAT3/g- globin/HbF	131
miR-494-3p	SOCS6	Up regulated	Indirectly via SOCS6 inhibiting	PMF	K562/cord blood CD34 ⁺ cells	miR-494-3p up regulated STAT3 phosphorylation via SOCS6 inhibiting	miR-494-3p/SOCS6/ STAT3	144
miR-23a, -27a, -24 cluster	miR-23a, -27a and -24 synergistically target the members of gp130/JAK1/ STAT3 signaling pathway	Down regulated	Directly increase in STAT3	AEL (AML M6)	HEL and K562 cell lines/CB-HSCS CD34 ⁺	miR-23a, -27a, -24 cluster down regulated STAT3	miR-23a, -27a, -24/ GP130/JAK1/STAT3 pathway	152

miRNA-155	N/A	Up regulated in some subtypes of AML	Indirectly via SOCS1	AML	HL60/Dami and THP1 cell lines	pSTAT3 down regulated miR-155	IL6 family/LiFRa/ miRNA-155/SOCS1/ STAT3	155
miR-17/miR-20a	P21/STAT3	Down regulated by HIF1a	Directly targeting STAT3	AML	U937T/NB4 cells	miR-17/miR-20a down regulated STAT3	HIF1a/CEBPa/c-Myc/ miR-17/miR-20a/ JAK/STAT signaling pathways	175

promoted cell cycle arrest and differentiation in myeloid leukemic cells.¹⁷⁵ The HIF-1a protein translocates into the nucleus and forms a heterodimer with HIF-1b and then regulates the expression of target genes through binding to hypoxia-responsive elements (HREs) located on gene promoters.¹⁷⁵ These genes, which are targeted by HIF-1, support the cells for adaptation in hypoxic conditions by affecting processes covering, apoptosis, differentiation, angiogenesis, cell growth in concomitant with metabolism, and erythropoiesis.¹⁷⁵ HIF-1a in hypoxic conditions, known as a significant indicator of solid tumors, cooperates with tumor growth, metastasis, and angiogenesis.¹⁷² He et al have shown that the HIF-1a transcription factor can induce differentiation and inhibit AML development.^{173,175–179} They showed that HIF1a decreases miR-17 and miR-20a, two members of the miR-17-92 gene cluster, via directly targeting STAT3.¹⁷⁵ It has already been found that miR-17 and miR-20a are overexpressed in solid tumors and hematological disorders such as mantle cell lymphoma (MCL), large B-cell lymphoma, and Burkitt’s lymphoma.^{180–182} Besides, it has been shown that the miR-17-92 cluster target the HIF-1a protein.^{175,183–185} He et al also reported that miR-17 and miR-20a were downregulated in hypoxic conditions in AML cell lines. Indeed, they indicated that HIF-1a downregulates miR-17 and miR-20a gene expression during hypoxia in AML cells. They also demonstrated that a decrease in miR-17 and miR-20a participates in the differentiation process of AML cell lines triggered by HIF-1a. Based on last year’s reports, STAT3 can be a target gene of miR-17 and miR-20a.^{186,187} On the other hand, they reported that exist two binding sites for miR-17/miR-20a at the wild-type 3’-UTR of STAT3, which enables suppressing of the STAT3 protein expression via directly targeting its 3’-UTR post-transcriptionally.¹⁷⁵ In total, He et al concluded that miR-17 or miR-20a abrogated HIF-1a-induced growth arrest and differentiation in AML cells through binding to STAT3 transcript and inhibiting its expression.¹⁷⁵

The miR-21 and miR-17-92 cluster and their association with STAT3

Acute myeloid leukemia (AML) with t(8;16)(p11;p13) is an uncommon leukemia subtype with representative clinical features, including presentation as coagulation disorder, recurrent extramedullary involvement accompanied by poor prognosis.¹⁸⁸ Further, AML blast cells with t(8;16) show a frequent hemophagocytosis and myelomonocytic phase of differentiation.^{188,189} The oncogenic miRNAs cluster miR-17-92 and miR-21, which are usually upregulated in other cancers, are commonly downregulated in the t(8;16) AML 191. In this regard, assessment of the expression levels of the known transcription factors of cluster miR-17-92 and miR-21 genes, such as STAT3 in 7 t(8;16) AML patients and 36 patients with other AML cytogenetic subtypes approved STAT3 downregulation in the t(8;16) AML patients 191. In 2013, Beya et al found that STAT3 expression was down-regulated in patients with t(8;16) translocation, and miR-21 was regulated at the transcriptional level by STAT3.¹⁸⁹ Based on these results, the decrease in miR-21 and likely decrease in miR-17-92 cluster has an association with a decrease in STAT3 in AML with t(8;16).¹⁸⁹

Conclusion

miRNAs regulate a variety of cellular processes, including inflammation, proliferation, survival, metastasis, invasion, and angiogenesis, all of which can also be triggered by activated STAT3 by binding to DNA and other stimulants, leading to tumor growth (Fig. 2, 3). Furthermore, the important role of miRNAs in regulating the JAK/STAT3 signaling pathway has been proven. For a better understanding of the interaction between STAT3 and miRNAs in leukemic cells, the complete signaling pathways have been shown in Figure 4. To determine the role of STAT3, one of the most important transcription factors in many cancers, including leukemia, we focused on studies regarding the role of miRNAs in the regulation JAK/STAT3 signaling pathway in hematological disorders. For a better understanding of the relationship between microRNAs and STAT3, here we have summarized the role of both factors in some cancers and more comprehensively in blood malignancies in two separate tables (Table 1, 2). To sum up, it can be concluded that therapeutic approaches especially multimodal treatments for modifying the STAT3 signaling pathway can lead to improved therapeutic outcomes in patients suffering from various hematological disorders, in particular, leukemia; however, the accomplishment of comprehensive clinical trials before using STATs inhibitors should be considered.

Author contributions

All authors contributed to the conception and the main idea of the work. M.S.D, T.A.M.M, H.S.R, and M.S.M drafted the main text, figures, and tables. S.D, N.S and S.S took part in the conceptualization and scientific editing of the manuscript. F.M and E.Z drew the figures. M.A, A.A and R.A helped in revision. M.J and M.F.H supervised the work and provided the comments and additional scientific information. All authors read and approved the final version of the work to be published.

Consent for publication

All authors consent to the publication of the manuscript in Genes & Diseases.

Conflict of interests

Authors declare no conflict of interests.

Acknowledgements

Authors thank the personnel of Immunology Research Center and Department of Hematology affiliated to Tabriz University of Medical Sciences.

References

- Mrózek K, Heerema NA, Bloomfield CD. Cytogenetics in acute leukemia. *Blood Rev.* 2004;18(2):115–136.
- Eisfeld AK, Kohlschmidt J, Mims A, et al. Additional gene mutations may refine the 2017 European LeukemiaNet classification in adult patients with de novo acute myeloid leukemia aged <60 years. *Leukemia.* 2020;34(12):3215–3227.
- Brown AL, Hahn CN, Scott HS. Secondary leukemia in patients with germline transcription factor mutations (RUNX1, GATA2, CEBPA). *Blood.* 2020;136(1):24–35.
- Ashihara E, Takada T, Maekawa T. Targeting the canonical Wnt/ β -catenin pathway in hematological malignancies. *Cancer Sci.* 2015;106(6):665–671.
- Tasian SK, Teachey DT, Rheingold SR. Targeting the PI3K/mTOR pathway in pediatric hematologic malignancies. *Front Oncol.* 2014;4:108.
- Ge X, Wang X. Role of Wnt canonical pathway in hematological malignancies. *J Hematol Oncol.* 2010;3:33.
- Vainchenker W, Constantinescu SN. JAK/STAT signaling in hematological malignancies. *Oncogene.* 2013;32(21):2601–2613.
- Xin P, Xu X, Deng C, et al. The role of JAK/STAT signaling pathway and its inhibitors in diseases. *Int Immunopharmacol.* 2020;80:106210.
- Pencik J, Pham HTT, Schmoellerl J, et al. JAK-STAT signaling in cancer: from cytokines to non-coding genome. *Cytokine.* 2016;87:26–36.
- Kleppe M, Koche R, Zou L, et al. Dual targeting of oncogenic activation and inflammatory signaling increases therapeutic efficacy in myeloproliferative neoplasms. *Cancer Cell.* 2018;33(1):29–43.
- Tolomeo M, Cascio A. The multifaced role of STAT3 in cancer and its implication for anticancer therapy. *Int J Mol Sci.* 2021;22(2):603.
- Constantinescu SN, Girardot M, Pecquet C. Mining for JAK–STAT mutations in cancer. *Trends Biochem Sci.* 2008;33(3):122–131.
- Arora L, Kumar AP, Arfuso F, Chng WJ, Sethi G. The role of signal transducer and activator of transcription 3 (STAT3) and its targeted inhibition in hematological malignancies. *Cancers (Basel).* 2018;10(9):327.
- Wu Z, Liu J, Hu S, Zhu Y, Li S. Serine/threonine kinase 35, a target gene of STAT3, regulates the proliferation and apoptosis of osteosarcoma cells. *Cell Physiol Biochem.* 2018;45(2):808–818.
- Boudny M, Trbusek M. The important role of STAT3 in chronic lymphocytic leukaemia biology. *Klin Onkol.* 2020;33(1):32–38.
- Nairismägi M-, Gerritsen ME, Li ZM, et al. Oncogenic activation of JAK3-STAT signaling confers clinical sensitivity to PRN371, a novel selective and potent JAK3 inhibitor, in natural killer/T-cell lymphoma. *Leukemia.* 2018;32(5):1147–1156.
- Thomas SJ, Snowden JA, Zeidler MP, Danson SJ. The role of JAK/STAT signalling in the pathogenesis, prognosis and treatment of solid tumours. *Br J Cancer.* 2015;113(3):365–371.
- Bose S, Banerjee S, Mondal A, et al. Targeting the JAK/STAT signaling pathway using phytocompounds for cancer prevention and therapy. *Cells.* 2020;9(6):1451.
- Benekli M, Baer MR, Baumann H, Wetzler M. Signal transducer and activator of transcription proteins in leukemias. *Blood.* 2003;101(8):2940–2954.

20. Zheng Q, Dong H, Mo J, et al. A novel STAT3 inhibitor W2014-S regresses human non-small cell lung cancer xenografts and sensitizes EGFR-TKI acquired resistance. *Theranostics*. 2021; 11(2):824–840.
21. Li P, Grgurevic S, Liu Z, et al. Signal transducer and activator of transcription–3 induces microRNA-155 expression in chronic lymphocytic leukemia. *PLoS One*. 2013;8(6): e64678.
22. Kuusanmäki H, Dufva O, Parri E, et al. Drug sensitivity profiling identifies potential therapies for lymphoproliferative disorders with overactive JAK/STAT3 signaling. *Oncotarget*. 2017;8(57):97516–97527.
23. Furtak SL, Backos DS, Matheson CJ, Reigan P. Strategies and approaches of targeting STAT3 for cancer treatment. *ACS Chem Biol*. 2016;11(2):308–318.
24. Zou S, Tong Q, Liu B, Huang W, Tian Y, Fu X. Targeting STAT3 in cancer immunotherapy. *Mol Cancer*. 2020;19(1):145.
25. Cao Q, Li YY, He WF, et al. Interplay between microRNAs and the STAT3 signaling pathway in human cancers. *Physiol Genomics*. 2013;45(24):1206–1214.
26. Shan D, Shang Y, Hu T. MicroRNA-411 inhibits cervical cancer progression by directly targeting STAT3. *Oncol Rep*. 2019; 27(3):349–358.
27. Xu S, Zhao N, Hui L, Song M, Miao ZW, Jiang XJ. MicroRNA-124-3p inhibits the growth and metastasis of nasopharyngeal carcinoma cells by targeting STAT3. *Oncol Rep*. 2016;35(3): 1385–1394.
28. Tian K, Liu W, Zhang J, et al. MicroRNA-125b exerts antitumor functions in cutaneous squamous cell carcinoma by targeting the STAT3 pathway. *Cell Mol Biol Lett*. 2020;25:12.
29. Cai K, Li HX, Li PP, Guo ZJ, Yang Y. MicroRNA-449b-3p inhibits epithelial-mesenchymal transition by targeting IL-6 and through the JAK2/STAT3 signaling pathway in non-small cell lung cancer. *Exp Ther Med*. 2020;19(4):2527–2534.
30. Wang B, Hsu SH, Frankel W, Ghoshal K, Jacob ST. Stat3-mediated activation of microRNA-23a suppresses gluconeogenesis in hepatocellular carcinoma by down-regulating glucose-6-phosphatase and peroxisome proliferator-activated receptor gamma, coactivator 1 alpha. *Hepatology*. 2012; 56(1):186–197.
31. Tang Z, Xu T, Li Y, Fei W, Yang G, Hong Y. Inhibition of CRY2 by STAT3/miRNA-7-5p promotes osteoblast differentiation through upregulation of CLOCK/BMAL1/P300 expression. *Mol Ther Nucleic Acids*. 2020;19:865–876.
32. Wang A, Deng S, Chen X, et al. miR-29a-5p/STAT3 positive feedback loop regulates TETs in colitis-associated colorectal cancer. *Inflamm Bowel Dis*. 2020;26(4):524–533.
33. Cheng M, Wang B, Yang M, et al. microRNAs expression in relation to particulate matter exposure: a systematic review. *Environ Pollut*. 2020;260:113961.
34. Azar MRMH, Akbari M, Mohammed HN, Asadi M, Mahdavi F. Dysregulation of miR-27a and SMAD2 can be a reliable indicator in the prognosis and diagnosis of CRC as well as in response to chemotherapy drugs. *Gene Rep*. 2020;21(3):100844.
35. Azar MRMH, Aghazadeh H, Mohammed HN, et al. miR-193a-5p as a promising therapeutic candidate in colorectal cancer by reducing 5-FU and Oxaliplatin chemoresistance by targeting CXCR4. *Int Immunopharmacol*. 2021;92:107355.
36. Liu D, Zhang N, Zhang X, Qin M, Dong Y, Jin L. MiR-410 down-regulates the expression of interleukin-10 by targeting STAT3 in the pathogenesis of systemic lupus erythematosus. *Cell Physiol Biochem*. 2016;39(1):303–315.
37. Niu L, Yang W, Duan L, et al. Biological implications and clinical potential of metastasis-related miRNA in colorectal cancer. *Mol Ther Nucleic Acids*. 2020;23:42–54.
38. He B, Zhao Z, Cai Q, et al. miRNA-based biomarkers, therapies, and resistance in cancer. *Int J Biol Sci*. 2020;16(14): 2628–2647.
39. Kabekkodu SP, Shukla V, Varghese VK, et al. Cluster miRNAs and cancer: diagnostic, prognostic and therapeutic opportunities. *Wiley Interdiscip Rev RNA*. 2020;11(2):e1563.
40. Tamjidifar R, Akbari M, Tarzi S, et al. Prognostic and diagnostic values of miR-506 and SPON 1 in colorectal cancer with clinicopathological considerations. *J Gastrointest Cancer*. 2021;52(1):125–129.
41. Rupaimoole R, Calin GA, Lopez-Berestein G, Sood AK. miRNA deregulation in cancer cells and the tumor microenvironment. *Cancer Discov*. 2016;6(3):235–246.
42. Raza U, Zhang JD, Sahin O. MicroRNAs: master regulators of drug resistance, stemness, and metastasis. *J Mol Med (Berl)*. 2014;92(4):321–336.
43. Nabipoorashrafi SA, Shomali N, Sadat-Hatamnezhad L, et al. miR-143 acts as an inhibitor of migration and proliferation as well as an inducer of apoptosis in melanoma cancer cells in vitro. *IUBMB Life*. 2020;72(9):2034–2044.
44. Samidurai A, Roh SK, Prakash M, et al. STAT3-miR-17/20 signalling axis plays a critical role in attenuating myocardial infarction following rapamycin treatment in diabetic mice. *Cardiovasc Res*. 2020;116(13):2103–2115.
45. Dong L, Cao X, Luo Y, Zhang G, Zhang D. A positive feedback loop of lncRNA DSCR8/miR-98-5p/STAT3/HIF-1 α plays a role in the progression of ovarian cancer. *Front Oncol*. 2020;10: 1713.
46. Li Z, Song Y, He T, et al. M2 microglial small extracellular vesicles reduce glial scar formation via the miR-124/STAT3 pathway after ischemic stroke in mice. *Theranostics*. 2021; 11(3):1232–1248.
47. Yan X, Zeng D, Zhu H, et al. MiRNA-532-5p regulates CUMS-induced depression-like behaviors and modulates LPS-induced proinflammatory cytokine signaling by targeting STAT3. *Neuropsychiatr Dis Treat*. 2020;16:2753–2764.
48. Chen N, Feng L, Qu H, et al. Overexpression of IL-9 induced by STAT3 phosphorylation is mediated by miR-155 and miR-21 in chronic lymphocytic leukemia. *Oncol Rep*. 2018;39(6): 3064–3072.
49. Carabia J, Carpio C, Abrisqueta P, et al. Microenvironment regulates the expression of miR-21 and tumor suppressor genes PTEN, PIAS3 and PDCD4 through ZAP-70 in chronic lymphocytic leukemia. *Sci Rep*. 2017;7(1):12262.
50. Soroosh P, Doherty TA. Th9 and allergic disease. *Immunology*. 2009;127(4):450–458.
51. Tete S, Saggini A, Maccauro G, et al. Interleukin-9 and mast cells. *J Biol Regul Homeost Agents*. 2012;26(3):319–326.
52. Hornakova T, Staerk J, Royer Y, et al. Acute lymphoblastic leukemia-associated JAK1 mutants activate the Janus kinase/STAT pathway via interleukin-9 receptor alpha homodimers. *J Biol Chem*. 2009;284(11):6773–6781.
53. Carretero R, Wang E, Rodriguez AI, et al. Regression of melanoma metastases after immunotherapy is associated with activation of antigen presentation and interferon-mediated rejection genes. *Int J Cancer*. 2012;131(2):387–395.
54. Randhawa J, Ostojic A, Vrhovac R, Atallah E, Verstovsek S. Splenomegaly in myelofibrosis—new options for therapy and the therapeutic potential of Janus kinase 2 inhibitors. *J Hematol Oncol*. 2012;5:43.
55. Byrd JC, Furman RR, Coutre SE, et al. Three-year follow-up of treatment-naïve and previously treated patients with CLL and SLL receiving single-agent ibrutinib. *Blood*. 2015;125(16): 2497–2506.
56. Calin GA, Ferracin M, Cimmino A, et al. A MicroRNA signature associated with prognosis and progression in chronic lymphocytic leukemia. *N Engl J Med*. 2005;353(17):1793–1801.
57. Saleh LM, Wang W, Herman SE, et al. Ibrutinib downregulates a subset of miRNA leading to upregulation of tumor suppressors and inhibition of cell proliferation in chronic lymphocytic leukemia. *Leukemia*. 2017;31(2):340–349.

58. Rossi S, Shimizu M, Barbarotto E, et al. microRNA fingerprinting of CLL patients with chromosome 17p deletion identify a miR-21 score that stratifies early survival. *Blood*. 2010;116(6):945–952.
59. Fu X, Han Y, Wu Y, et al. Prognostic role of microRNA-21 in various carcinomas: a systematic review and meta-analysis. *Eur J Clin Invest*. 2011;41(11):1245–1253.
60. Jazbutyte V, Thum T. MicroRNA-21: from cancer to cardiovascular disease. *Curr Drug Targets*. 2010;11(8):926–935.
61. Xiong Q, Zhong Q, Zhang J, et al. Identification of novel miR-21 target proteins in multiple myeloma cells by quantitative proteomics. *J Proteome Res*. 2012;11(4):2078–2090.
62. Frankel LB, Christoffersen NR, Jacobsen A, Lindow M, Krogh L, Lund AH. Programmed cell death 4 (PDCD4) is an important functional target of the microRNA miR-21 in breast cancer cells. *J Biol Chem*. 2008;283(2):1026–1033.
63. Wickramasinghe NS, Manavalan TT, Dougherty SM, Riggs KA, Li Y, Klinge CM. Estradiol downregulates miR-21 expression and increases miR-21 target gene expression in MCF-7 breast cancer cells. *Nucleic Acids Res*. 2009;37(8):2584–2595.
64. Löffler D, Brocke-Heidrich K, Pfeifer G, et al. Interleukin-6 dependent survival of multiple myeloma cells involves the Stat3-mediated induction of microRNA-21 through a highly conserved enhancer. *Blood*. 2007;110(4):1330–1333.
65. Leng RX, Pan HF, Qin WZ, Chen GM, Ye DQ. Role of microRNA-155 in autoimmunity. *Cytokine Growth Factor Rev*. 2011;22(3):141–147.
66. Costinean S, Zanasi N, Pekarsky Y, et al. Pre-B cell proliferation and lymphoblastic leukemia/high-grade lymphoma in E(mu)-miR155 transgenic mice. *Proc Natl Acad Sci U S A*. 2006;103(18):7024–7029.
67. Thai TH, Calado DP, Casola S, et al. Regulation of the germinal center response by microRNA-155. *Science*. 2007;316(5824):604–608.
68. Haasch D, Chen YW, Reilly RM, et al. T cell activation induces a noncoding RNA transcript sensitive to inhibition by immunosuppressant drugs and encoded by the proto-oncogene, BIC. *Cell Immunol*. 2002;217(1–2):78–86.
69. Eis PS, Tam W, Sun L, et al. Accumulation of miR-155 and BIC RNA in human B cell lymphomas. *Proc Natl Acad Sci U S A*. 2005;102(10):3627–3632.
70. Fulci V, Chiaretti S, Goldoni M, et al. Quantitative technologies establish a novel microRNA profile of chronic lymphocytic leukemia. *Blood*. 2007;109(11):4944–4951.
71. Kluiver J, Poppema S, de Jong D, et al. BIC and miR-155 are highly expressed in Hodgkin, primary mediastinal and diffuse large B cell lymphomas. *J Pathol*. 2005;207(2):243–249.
72. Metzler M, Wilda M, Busch K, Viehmann S, Borkhardt A. High expression of precursor microRNA-155/BIC RNA in children with Burkitt lymphoma. *Genes Chromosomes Cancer*. 2004;39(2):167–169.
73. Hazan-Halevy I, Harris D, Liu Z, et al. STAT3 is constitutively phosphorylated on serine 727 residues, binds DNA, and activates transcription in CLL cells. *Blood*. 2010;115(14):2852–2863.
74. Li P, Harris D, Liu Z, Liu J, Keating M, Estrov Z. Stat3 activates the receptor tyrosine kinase like orphan receptor-1 gene in chronic lymphocytic leukemia cells. *PLoS One*. 2010;5(7):e11859.
75. Tivnan A, Tracey L, Buckley PG, Alcock LC, Davidoff AM, Stallings RL. MicroRNA-34a is a potent tumor suppressor molecule in vivo in neuroblastoma. *BMC Cancer*. 2011;11:33.
76. Vargova K, Curik N, Burda P, et al. MYB transcriptionally regulates the miR-155 host gene in chronic lymphocytic leukemia. *Blood*. 2011;117(14):3816–3825.
77. Aaronson DS, Horvath CM. A road map for those who don't know JAK-STAT. *Science*. 2002;296(5573):1653–1655.
78. Marinescu C, Vlădăreanu AM, Mihai F. Acute lymphocytic leukemia in adults. Pathologic features and prognosis. *Rom J Intern Med*. 2015;53(1):31–36.
79. Den Boer ML, van Slegtenhorst M, De Menezes RX, et al. A subtype of childhood acute lymphoblastic leukaemia with poor treatment outcome: a genome-wide classification study. *Lancet Oncol*. 2009;10(2):125–134.
80. Igwe JJ, Yang D, Merchant A, et al. The presence of Philadelphia chromosome does not confer poor prognosis in adult pre-B acute lymphoblastic leukaemia in the tyrosine kinase inhibitor era – a surveillance, epidemiology, and end results database analysis. *Br J Haematol*. 2017;179(4):618–626.
81. Möricke A, Reiter A, Zimmermann M, et al. Risk-adjusted therapy of acute lymphoblastic leukemia can decrease treatment burden and improve survival: treatment results of 2169 unselected pediatric and adolescent patients enrolled in the trial ALL-BFM 95. *Blood*. 2008;111(9):4477–4489.
82. Reynaud D, Pietras E, Barry-Holmes K, et al. IL-6 controls leukemic multipotent progenitor cell fate and contributes to chronic myelogenous leukemia development. *Cancer Cell*. 2011;20(5):661–673.
83. Jiang T, Chen J, Huang XB, Li YX, Zhong L. miR-451a induced apoptosis of Philadelphia chromosome-positive acute lymphoblastic leukemia cells by targeting IL-6R. *Neoplasia*. 2018;65(6):907–914.
84. Wang SW, Sun YM. The IL-6/JAK/STAT3 pathway: potential therapeutic strategies in treating colorectal cancer (Review). *Int J Oncol*. 2014;44(4):1032–1040.
85. Garbers C, Aparicio-Siegmund S, Rose-John S. The IL-6/gp130/STAT3 signaling axis: recent advances towards specific inhibition. *Curr Opin Immunol*. 2015;34:75–82.
86. Soltani I, Douzi K, Gharbi H, et al. Downregulation of miR-451 in Tunisian chronic myeloid leukemia patients: potential implication in imatinib resistance. *Hematology*. 2017;22(4):201–207.
87. Chen Q, Hu H, Jiao D, et al. miR-126-3p and miR-451a correlate with clinicopathological features of lung adenocarcinoma: the underlying molecular mechanisms. *Oncol Rep*. 2016;36(2):909–917.
88. Su Z, Ni L, Yu W, et al. MicroRNA-451a is associated with cell proliferation, migration and apoptosis in renal cell carcinoma. *Mol Med Rep*. 2015;11(3):2248–2254.
89. Lopotová T, Žáčková M, Klamová H, Moravcová J. MicroRNA-451 in chronic myeloid leukemia: miR-451–BCR-ABL regulatory loop? *Leuk Res*. 2011;35(7):974–977.
90. Lamy T, Moignot A, Loughran Jr TP. LGL leukemia: from pathogenesis to treatment. *Blood*. 2017;129(9):1082–1094.
91. Semenzato G, Zambello R, Starkebaum G, Oshimi K, Loughran Jr TP. The lymphoproliferative disease of granular lymphocytes: updated criteria for diagnosis. *Blood*. 1997;89(1):256–260.
92. Mariotti B, Calabretto G, Rossato M, et al. Identification of a miR-146b-Fas ligand axis in the development of neutropenia in T large granular lymphocyte leukemia. *Haematologica*. 2020;105(5):1351–1360.
93. Leblanc F, Zhang D, Liu X, Loughran TP. Large granular lymphocyte leukemia: from dysregulated pathways to therapeutic targets. *Future Oncol*. 2012;8(7):787–801.
94. Epling-Burnette PK, Zhong B, Bai F, et al. Cooperative regulation of Mcl-1 by Janus kinase/stat and phosphatidylinositol 3-kinase contribute to granulocyte-macrophage colony-stimulating factor-delayed apoptosis in human neutrophils. *J Immunol*. 2001;166(12):7486–7495.

95. Teramo A, Barilà G, Calabretto G, et al. STAT3 mutation impacts biological and clinical features of T-LGL leukemia. *Oncotarget*. 2017;8(37):61876–61889.
96. Tanaka M, Suda T, Haze K, et al. Fas ligand in human serum. *Nat Med*. 1996;2(3):317–322.
97. Saitoh T, Karasawa M, Sakuraya M, et al. Improvement of extrathymic T cell type of large granular lymphocyte (LGL) leukemia by cyclosporin A: the serum level of Fas ligand is a marker of LGL leukemia activity. *Eur J Haematol*. 2000;65(4):272–275.
98. Liu JH, Wei S, Lamy T, et al. Chronic neutropenia mediated by fas ligand. *Blood*. 2000;95(10):3219–3222.
99. Perzova R, Loughran Jr TP. Constitutive expression of Fas ligand in large granular lymphocyte leukaemia. *Br J Haematol*. 1997;97(1):123–126.
100. Jerez A, Clemente MJ, Makishima H, et al. STAT3 mutations unify the pathogenesis of chronic lymphoproliferative disorders of NK cells and T-cell large granular lymphocyte leukemia. *Blood*. 2012;120(15):3048–3057.
101. Koskela HL, Eldfors S, Ellonen P, et al. Somatic STAT3 mutations in large granular lymphocytic leukemia. *N Engl J Med*. 2012;366(20):1905–1913.
102. Lee H, Deng J, Xin H, Liu Y, Pardoll D, Yu H. A requirement of STAT3 DNA binding precludes Th-1 immunostimulatory gene expression by NF- κ B in tumors. *Cancer Res*. 2011;71(11):3772–3780.
103. Niu G, Wright KL, Ma Y, et al. Role of Stat3 in regulating p53 expression and function. *Mol Cell Biol*. 2005;25(17):7432–7440.
104. Xiang M, Birkbak NJ, Vafaizadeh V, et al. STAT3 induction of miR-146b forms a feedback loop to inhibit the NF- κ B to IL-6 signaling axis and STAT3-driven cancer phenotypes. *Sci Signal*. 2014;7(310):ra11.
105. Vilella D, Ramalho RF, Silva AR, et al. Differential DNA methylation of microRNA genes in temporal cortex from Alzheimer's disease individuals. *Neural Plast*. 2016;2016:2584940.
106. Curtale G, Mirolo M, Renzi TA, Rossato M, Bazzoni F, Locati M. Negative regulation of Toll-like receptor 4 signaling by IL-10-dependent microRNA-146b. *Proc Natl Acad Sci U S A*. 2013;110(28):11499–11504.
107. Renzi TA, Rubino M, Gornati L, Garlanda C, Locati M, Curtale G. MiR-146b mediates endotoxin tolerance in human phagocytes. *Mediators Inflamm*. 2015;2015:145305.
108. Cheng HS, Sivachandran N, Lau A, et al. Micro RNA-146 represses endothelial activation by inhibiting pro-inflammatory pathways. *EMBO Mol Med*. 2013;5(7):1017–1034.
109. Drury GL, Di Marco S, Dormoy-Raclet V, Desbarats J, Gallouzi IE. FasL expression in activated T lymphocytes involves HuR-mediated stabilization. *J Biol Chem*. 2010;285(41):31130–31138.
110. Yang W, Yu H, Shen Y, Liu Y, Yang Z, Sun T. MiR-146b-5p overexpression attenuates stemness and radioresistance of glioma stem cells by targeting HuR/lincRNA-p21/ β -catenin pathway. *Oncotarget*. 2016;7(27):41505–41526.
111. Jabbour E, Kantarjian H. Chronic myeloid leukemia: 2014 update on diagnosis, monitoring, and management. *Am J Hematol*. 2014;89(5):547–556.
112. Zhang Y, Zhang HE, Liu Z. MicroRNA-147 suppresses proliferation, invasion and migration through the AKT/mTOR signaling pathway in breast cancer. *Oncol Lett*. 2016;11(1):405–410.
113. Sui CJ, Xu F, Shen WF, et al. MicroRNA-147 suppresses human hepatocellular carcinoma proliferation migration and chemosensitivity by inhibiting HOXC6. *Am J Cancer Res*. 2016;6(12):2787–2798.
114. Lee CG, McCarthy S, Gruidl M, Timme C, Yeatman TJ. MicroRNA-147 induces a mesenchymal-to-epithelial transition (MET) and reverses EGFR inhibitor resistance. *PLoS One*. 2014;9(1):e84597.
115. Chu G, Zhang J, Chen X. Serum level of microRNA-147 as diagnostic biomarker in human non-small cell lung cancer. *J Drug Target*. 2016;24(7):613–617.
116. Yao Y, Suo AL, Li ZF, et al. MicroRNA profiling of human gastric cancer. *Mol Med Rep*. 2009;2(6):963–970.
117. Han L, Dong Z, Liu N, Xie F, Wang N. Maternally expressed gene 3 (MEG3) enhances PC12 cell hypoxia injury by targeting MiR-147. *Cell Physiol Biochem*. 2017;43(6):2457–2469.
118. Li ZY, Yang L, Liu XJ, Wang XZ, Pan YX, Luo JM. The long noncoding RNA MEG3 and its target miR-147 regulate JAK/STAT pathway in advanced chronic myeloid leukemia. *EBio-Medicine*. 2018;34:61–75.
119. Tatarano S, Chiyomaru T, Kawakami K, et al. Novel oncogenic function of mesoderm development candidate 1 and its regulation by MiR-574-3p in bladder cancer cell lines. *Int J Oncol*. 2012;40(4):951–959.
120. Krishnan P, Ghosh S, Wang B, et al. Next generation sequencing profiling identifies miR-574-3p and miR-660-5p as potential novel prognostic markers for breast cancer. *BMC Genomics*. 2015;16(1):735.
121. Su Y, Ni Z, Wang G, et al. Aberrant expression of microRNAs in gastric cancer and biological significance of miR-574-3p. *Int Immunopharmacol*. 2012;13(4):468–475.
122. Xu H, Liu X, Zhou J, Chen X, Zhao J. miR-574-3p acts as a tumor promoter in osteosarcoma by targeting SMAD4 signaling pathway. *Oncol Lett*. 2016;12(6):5247–5253.
123. Okumura T, Kojima H, Miwa T, et al. The expression of microRNA 574-3p as a predictor of postoperative outcome in patients with esophageal squamous cell carcinoma. *World J Surg Oncol*. 2016;14(1):228.
124. Hong DS, Angelo LS, Kurzrock R. Interleukin-6 and its receptor in cancer: implications for translational therapeutics. *Cancer*. 2007;110(9):1911–1928.
125. Maeda K, Baba Y, Nagai Y, et al. IL-6 blocks a discrete early step in lymphopoiesis. *Blood*. 2005;106(3):879–885.
126. Ciarcia R, Vitiello MT, Galdiero M, et al. Imatinib treatment inhibit IL-6, IL-8, NF-KB and AP-1 production and modulate intracellular calcium in CML patients. *J Cell Physiol*. 2012;227(6):2798–2803.
127. Yang H, Zhang J, Li J, et al. Overexpression of miR-574-3p suppresses proliferation and induces apoptosis of chronic myeloid leukemia cells via targeting IL6/JAK/STAT3 pathway. *Exp Ther Med*. 2018;16(5):4296–4302.
128. Piel FB, Patil AP, Howes RE, et al. Global epidemiology of sickle haemoglobin in neonates: a contemporary geo-statistical model-based map and population estimates. *Lancet*. 2013;381(9861):142–151.
129. Dong M, McGann PT. Changing the clinical paradigm of hydroxyurea treatment for sickle cell anemia through precision medicine. *Clin Pharmacol Ther*. 2021;109(1):73–81.
130. Platt OS, Brambilla DJ, Rosse WF, et al. Mortality in sickle cell disease. Life expectancy and risk factors for early death. *N Engl J Med*. 1994;330(23):1639–1644.
131. Ward CM, Li B, Pace BS. Original Research: Stable expression of miR-34a mediates fetal hemoglobin induction in K562 cells. *Exp Biol Med (Maywood)*. 2016;241(7):719–729.
132. Poillon WN, Kim BC, Rodgers GP, Noguchi CT, Schechter AN. Sparing effect of hemoglobin F and hemoglobin A2 on the polymerization of hemoglobin S at physiologic ligand saturations. *Proc Natl Acad Sci U S A*. 1993;90(11):5039–5043.

133. Vekilov PG. Sick-cell haemoglobin polymerization: is it the primary pathogenic event of sickle-cell anaemia? *Br J Haematol.* 2007;139(2):173–184.
134. Weng W, Wang M, Xie S, et al. YY1-C/EBP α -miR34a regulatory circuitry is involved in renal cell carcinoma progression. *Oncol Rep.* 2014;31(4):1921–1927.
135. Chen QR, Yu LR, Tsang P, et al. Systematic proteome analysis identifies transcription factor YY1 as a direct target of miR-34a. *J Proteome Res.* 2011;10(2):479–487.
136. Zhao J, Lammers P, Torrance CJ, Bader AG. TP53-independent function of miR-34a via HDAC1 and p21(CIP1/WAF1.). *Mol Ther.* 2013;21(9):1678–1686.
137. Lal A, Thomas MP, Altschuler G, et al. Capture of microRNA-bound mRNAs identifies the tumor suppressor miR-34a as a regulator of growth factor signaling. *PLoS Genet.* 2011;7(11):e1002363.
138. Yao X, Kodeboyina S, Liu L, et al. Role of STAT3 and GATA-1 interactions in gamma-globin gene expression. *Exp Hematol.* 2009;37(8):889–900.
139. Wierenga AT, Vogelzang I, Eggen BJ, Vellenga E. Erythropoietin-induced serine 727 phosphorylation of STAT3 in erythroid cells is mediated by a MEK-, ERK-, and MSK1-dependent pathway. *Exp Hematol.* 2003;31(5):398–405.
140. De Kouchkovsky I, Abdul-Hay M. Acute myeloid leukemia: a comprehensive review and 2016 update. *Blood Cancer J.* 2016;6(7):e441.
141. Hitchcock IS, Kaushansky K. Thrombopoietin from beginning to end. *Br J Haematol.* 2014;165(2):259–268.
142. Drachman JG, Sabath DF, Fox NE, Kaushansky K. Thrombopoietin signal transduction in purified murine megakaryocytes. *Blood.* 1997;89(2):483–492.
143. Norfo R, Zini R, Pennucci V, et al. miRNA-mRNA integrative analysis in primary myelofibrosis CD34+ cells: role of miR-155/JARID2 axis in abnormal megakaryopoiesis. *Blood.* 2014;124(13):e21–e32.
144. Rontautori S, Norfo R, Pennucci V, et al. miR-494-3p overexpression promotes megakaryocytopoiesis in primary myelofibrosis hematopoietic stem/progenitor cells by targeting SOCS6. *Oncotarget.* 2017;8(13):21380–21397.
145. Shen R, Wang Y, Wang CX, et al. MiRNA-155 mediates TAM resistance by modulating SOCS6-STAT3 signalling pathway in breast cancer. *Am J Transl Res.* 2015;7(10):2115–2126.
146. Hwang MN, Min CH, Kim HS, et al. The nuclear localization of SOCS6 requires the N-terminal region and negatively regulates Stat3 protein levels. *Biochem Biophys Res Commun.* 2007;360(2):333–338.
147. Goldberg SL, Noel P, Klumpp TR, Dewald GW. The erythroid leukemias: a comparative study of erythroleukemia (FAB M6) and Di Guglielmo disease. *Am J Clin Oncol.* 1998;21(1):42–47.
148. Wells AW, Bown N, Reid MM, Hamilton PJ, Jackson GH, Taylor PR. Erythroleukaemia in the north of England: a population based study. *J Clin Pathol.* 2001;54(8):608–612.
149. Hasserjian RP, Zuo Z, Garcia C, et al. Acute erythroid leukemia: a reassessment using criteria refined in the 2008 WHO classification. *Blood.* 2010;115(10):1985–1992.
150. Santos FP, Faderl S, Garcia-Manero G, et al. Adult acute erythroleukemia: an analysis of 91 patients treated at a single institution. *Leukemia.* 2009;23(12):2275–2280.
151. Hegde S, Ni S, He S, et al. Stat3 promotes the development of erythroleukemia by inducing Pu. 1 expression and inhibiting erythroid differentiation. *Oncogene.* 2009;28(38):3349–3359.
152. Su R, Dong L, Zou D, et al. microRNA-23a, -27a and -24 synergistically regulate JAK1/Stat3 cascade and serve as novel therapeutic targets in human acute erythroid leukemia. *Oncogene.* 2016;35(46):6001–6014.
153. Zhao J, Xu Y, Zong Y, et al. Inhibition of Stat3 expression induces apoptosis and suppresses proliferation in human leukemia HL-60 cells. *Hematology.* 2011;16(4):232–235.
154. Epling-Burnette PK, Liu JH, Catlett-Falcone R, et al. Inhibition of STAT3 signaling leads to apoptosis of leukemic large granular lymphocytes and decreased Mcl-1 expression. *J Clin Invest.* 2001;107(3):351–362.
155. Xu S, Xu Z, Liu B, et al. LIFR α -CT3 induces differentiation of a human acute myelogenous leukemia cell line HL-60 by suppressing miR-155 expression through the JAK/STAT pathway. *Leuk Res.* 2014;38(10):1237–1244.
156. Guo SY, Shen X, Yang J, et al. TIMP-1 mediates the inhibitory effect of interleukin-6 on the proliferation of a hepatocarcinoma cell line in a STAT3-dependent manner. *Braz J Med Biol Res.* 2007;40(5):621–631.
157. Ying J, Tsujii M, Kondo J, et al. The effectiveness of an anti-human IL-6 receptor monoclonal antibody combined with chemotherapy to target colon cancer stem-like cells. *Int J Oncol.* 2015;46(4):1551–1559.
158. Cheng X, Jin G, Zhang X, Tian M, Zou L. Stage-dependent STAT3 activation is involved in the differentiation of rat hippocampus neural stem cells. *Neurosci Lett.* 2011;493(1–2):18–23.
159. Steyn PJ, Dzobo K, Smith RI, Myburgh KH. Interleukin-6 induces myogenic differentiation via JAK2-STAT3 signaling in mouse C2C12 myoblast cell line and primary human myoblasts. *Int J Mol Sci.* 2019;20(21):5273.
160. Nakajima K, Yamanaka Y, Nakae K, et al. A central role for Stat3 in IL-6-induced regulation of growth and differentiation in M1 leukemia cells. *EMBO J.* 1996;15(14):3651–3658.
161. Chonov DC, Ignatova MMK, Ananiev JR, Gulubova MV. IL-6 activities in the tumour microenvironment. Part 1. *Open Access Maced J Med Sci.* 2019;7(14):2391–2398.
162. Taga T, Kishimoto T. Gp130 and the interleukin-6 family of cytokines. *Annu Rev Immunol.* 1997;15:797–819.
163. Sun Q, Wang J, Xiong J, Yang L, Liu H. Free LIF receptor α -chain distal cytoplasmic motifs enhance Jak2-independent STAT3 phosphorylation and induce differentiation in HL-60 cells. *Oncol Rep.* 2011;26(2):399–404.
164. Lagos-Quintana M, Rauhut R, Yalcin A, Meyer J, Lendeckel W, Tuschl T. Identification of tissue-specific microRNAs from mouse. *Curr Biol.* 2002;12(9):735–739.
165. Jongen-Lavrencic M, Sun SM, Dijkstra MK, Valk PJ, Löwenberg B. MicroRNA expression profiling in relation to the genetic heterogeneity of acute myeloid leukemia. *Blood.* 2008;111(10):5078–5085.
166. Marcucci G, Mrózek K, Radmacher MD, Garzon R, Bloomfield CD. The prognostic and functional role of microRNAs in acute myeloid leukemia. *Blood.* 2011;117(4):1121–1129.
167. Garzon R, Volinia S, Liu CG, et al. MicroRNA signatures associated with cytogenetics and prognosis in acute myeloid leukemia. *Blood.* 2008;111(6):3183–3189.
168. McCoy CE, Sheedy FJ, Qualls JE, et al. IL-10 inhibits miR-155 induction by toll-like receptors. *J Biol Chem.* 2010;285(27):20492–20498.
169. Starr R, Willson TA, Viney EM, et al. A family of cytokine-inducible inhibitors of signalling. *Nature.* 1997;387(6636):917–921.
170. Semenza GL. Oxygen sensing, homeostasis, and disease. *N Engl J Med.* 2011;365(6):537–547.
171. Marofi F, Shomali N, Younus LA, et al. Under hypoxic conditions, MSCs affect the expression and methylation level of survival-related genes in ALL independent of apoptosis pathways in vitro. *Biotechnol Appl Biochem.* 2021. <https://doi.org/10.1002/bab.2154>.
172. Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell.* 2011;144(5):646–674.
173. Huang Y, Du KM, Xue ZH, et al. Cobalt chloride and low oxygen tension trigger differentiation of acute myeloid leukemic

- cells: possible mediation of hypoxia-inducible factor-1 α . *Leukemia*. 2003;17(11):2065–2073.
174. Zhao XY, Zhao KW, Jiang Y, Zhao M, Chen GQ. Synergistic induction of galectin-1 by CCAAT/enhancer binding protein α and hypoxia-inducible factor 1 α and its role in differentiation of acute myeloid leukemic cells. *J Biol Chem*. 2011;286(42):36808–36819.
 175. He M, Wang QY, Yin QQ, et al. HIF-1 α downregulates miR-17/20a directly targeting p21 and STAT3: a role in myeloid leukemic cell differentiation. *Cell Death Differ*. 2013;20(3):408–418.
 176. Kim JS, Cho EW, Chung HW, Kim IG. Effects of Tiron, 4, 5-dihydroxy-1, 3-benzene disulfonic acid, on human promyelocytic HL-60 leukemia cell differentiation and death. *Toxicology*. 2006;223(1–2):36–45.
 177. Liu W, Guo M, Xu YB, et al. Induction of tumor arrest and differentiation with prolonged survival by intermittent hypoxia in a mouse model of acute myeloid leukemia. *Blood*. 2006;107(2):698–707.
 178. Nguyen-Khac F, Della Valle V, Lopez RG, et al. Functional analyses of the TEL-ARNT fusion protein underscores a role for oxygen tension in hematopoietic cellular differentiation. *Oncogene*. 2006;25(35):4840–4847.
 179. Song LP, Zhang J, Wu SF, et al. Hypoxia-inducible factor-1 α -induced differentiation of myeloid leukemic cells is its transcriptional activity independent. *Oncogene*. 2008;27(4):519–527.
 180. Hayashita Y, Osada H, Tatematsu Y, et al. A polycistronic microRNA cluster, miR-17-92, is overexpressed in human lung cancers and enhances cell proliferation. *Cancer Res*. 2005;65(21):9628–9632.
 181. He L, Thomson JM, Hemann MT, et al. A microRNA polycistron as a potential human oncogene. *Nature*. 2005;435(7043):828–833.
 182. Volinia S, Calin GA, Liu CG, et al. A microRNA expression signature of human solid tumors defines cancer gene targets. *Proc Natl Acad Sci U S A*. 2006;103(7):2257–2261.
 183. Taguchi A, Yanagisawa K, Tanaka M, et al. Identification of hypoxia-inducible factor-1 α as a novel target for miR-17-92 microRNA cluster. *Cancer Res*. 2008;68(14):5540–5545.
 184. Ghosh G, Subramanian IV, Adhikari N, et al. Hypoxia-induced microRNA-424 expression in human endothelial cells regulates HIF- α isoforms and promotes angiogenesis. *J Clin Invest*. 2010;120(11):4141–4154.
 185. Cascio S, D'Andrea A, Ferla R, et al. miR-20b modulates VEGF expression by targeting HIF-1 α and STAT3 in MCF-7 breast cancer cells. *J Cell Physiol*. 2010;224(1):242–249.
 186. Abbas T, Dutta A. p21 in cancer: intricate networks and multiple activities. *Nat Rev Cancer*. 2009;9(6):400–414.
 187. Murray PJ. The JAK-STAT signaling pathway: input and output integration. *J Immunol*. 2007;178(5):2623–2629.
 188. Xie W, Hu S, Xu J, Chen Z, Medeiros LJ, Tang G. Acute myeloid leukemia with t (8; 16)(p11. 2; p13. 3)/KAT6A-CREBBP in adults. *Ann Hematol*. 2019;98(5):1149–1157.
 189. Díaz-Beyá M, Navarro A, Ferrer G, et al. Acute myeloid leukemia with translocation (8; 16)(p11; p13) and MYST3-CREBBP rearrangement harbors a distinctive microRNA signature targeting RET proto-oncogene. *Leukemia*. 2013;27(3):595–603.
 190. Park GB, Kim D. MicroRNA-503-5p inhibits the CD97-mediated JAK2/STAT3 pathway in metastatic or paclitaxel-resistant ovarian cancer cells. *Neoplasia*. 2019;21(2):206–215.
 191. Yu, Wu D, Gao H, et al. Clinical utility of a STAT3-regulated miRNA-200 family signature with prognostic potential in early gastric cancer. *Clin Cancer Res*. 2018;24(6):1459–1472.
 192. Chen MW, Yang ST, Chien MH, et al. The STAT3-miRNA-92-Wnt signaling pathway regulates spheroid formation and malignant progression in ovarian cancer. *Cancer Res*. 2017;77(8):1955–1967.
 193. Rokavec M, Öner MG, Li H, et al. IL-6R/STAT3/miR-34a feedback loop promotes EMT-mediated colorectal cancer invasion and metastasis. *J Clin Invest*. 2014;124(4):1853–1867.
 194. Hu H, Zhang Q, Chen W, et al. MicroRNA-301a promotes pancreatic cancer invasion and metastasis through the JAK/STAT3 signaling pathway by targeting SOCS5. *J Carcinog*. 2020;41(4):502–514.
 195. Wang X, Qiu W, Zhang G, et al. MicroRNA-204 targets JAK2 in breast cancer and induces cell apoptosis through the STAT3/BCI-2/survivin pathway. *Int J Clin Exp Pathol*. 2015;8(5):5017–5025.
 196. Zhou W, Bi X, Gao G, Sun L. miRNA-133b and miRNA-135a induce apoptosis via the JAK2/STAT3 signaling pathway in human renal carcinoma cells. *Biomed Pharmacother*. 2016;84:722–729.
 197. Patel K, Kollory A, Takashima A, Sarkar S, Faller DV, Ghosh SKJCL. MicroRNA let-7 downregulates STAT3 phosphorylation in pancreatic cancer cells by increasing SOCS3 expression. *Cancer Lett*. 2014;347(1):54–64.
 198. Zhang Y, Li X, Zhang J, Liang H. Natural killer T cell cytotoxic activity in cervical cancer is facilitated by the LINC00240/-microRNA-124-3p/STAT3/MICA axis. *Cancer Lett*. 2020;474:63–73.
 199. Haghikia A, Hoch M, Stapel B, Hilfiker-Kleiner D. STAT3 regulation of and by microRNAs in development and disease. *JAKSTAT*. 2012;1(3):143–150.
 200. El-Daly SM, Omara EA, Hussein J, Youness ER, El-Khayat ZJM. Differential expression of miRNAs regulating NF- κ B and STAT3 crosstalk during colitis-associated tumorigenesis. *Mol Cell Probes*. 2019;47:101442.
 201. Xue D, Yang Y., Liu Y, et al. MicroRNA-206 attenuates the growth and angiogenesis in non-small cell lung cancer cells by blocking the 14-3-3 ζ /STAT3/HIF-1 α /VEGF signaling. *Oncotarget*. 2016;7(48):79805–79813.
 202. Wei R, Yang Q, Han B, et al. microRNA-375 inhibits colorectal cancer cells proliferation by downregulating JAK2/STAT3 and MAP3K8/ERK signaling pathways. *Oncotarget*. 2017;8(10):16633–16641.
 203. Tao Y, Yang S, Wu Y, et al. MicroRNA-216a inhibits the metastasis of gastric cancer cells by targeting JAK2/STAT3-mediated EMT process. *Oncotarget*. 2017;8(51):88870–88881.
 204. Xu Y, Han YF, Zhu SJ, Dong JD, Ye B. miRNA-148a inhibits cell growth of papillary thyroid cancer through STAT3 and PI3K/AKT signaling pathways. *Oncol Rep*. 2017;38(5):3085–3093.
 205. Yang Y, Ding L, Hu Q, et al. MicroRNA-218 functions as a tumor suppressor in lung cancer by targeting IL-6/STAT3 and negatively correlates with poor prognosis. *Mol Cancer*. 2017;16(1):1–13.
 206. Mariotti B, Calabretto G, Rossato M, et al. Identification of a miR-146b-FasL axis in the development of neutropenia in T large granular lymphocyte leukemia. *Haematologica*. 2019;49:90.