



REVIEW ARTICLE

miR-106b as an emerging therapeutic target in cancer

Surendra Kumar Sagar

Department of Zoology, Swami Shraddhanand College, University of Delhi, Delhi 110036, India

Received 5 October 2020; received in revised form 24 January 2021; accepted 3 February 2021

Available online 12 February 2021

KEYWORDS

Apoptosis;
Cancer;
Metastases;
miRNA;
Oncogene;
Tumor suppressor

Abstract MicroRNAs (miRNAs) comprise short non-coding RNAs that function in regulating the expression of tumor suppressors or oncogenes and modulate oncogenic signaling pathways in cancer. miRNAs expression alters significantly in several tumor tissues and cancer cell lines. For example, miR-106b functions as an oncogene and increases in multiple cancers. The miR-106b directly targets genes involved in tumorigenesis, proliferation, invasion, migration, and metastases. This review has focused on the miR-106b function and its downstream target in different cancers and provide perspective into how miR-106 regulates cancer cell proliferation, migration, invasion, and metastases by regulating the tumor suppressor genes. Since miRNAs-based therapies are currently being developed to enhance cancer therapy outcomes, miR-106b could be an attractive and prospective candidate in different cancer types for detection, diagnosis, and prognosis assessment in the tumor.

Copyright © 2021, Chongqing Medical University. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

MicroRNAs (miRNAs) consists of short non-coding RNAs that control several gene expression. Recently, miRNAs have appeared as a potential molecular therapeutic approach for different diseases such as cancer.¹ In cancer cells, the aberrant miRNAs expression has been observed in different cancer and affects the hallmarks for tumor initiation and

progression. Several miRNAs play essential roles in the resistance of malignant cells to many anti-cancer agents.^{2,3} Most miRNAs are transcribed into primary miRNAs from DNA sequences and then processed into precursor miRNAs and mature miRNAs. Furthermore, miRNAs have been shown to bind with the target mRNAs to regulate the several genes involved in disease progression and cell death pathways.⁴ miRNAs behave either as a tumor suppressor or oncogene based on their targets.⁵

miR-106b overexpression has been stated in multiple tumor types and controls cell proliferation, migration, invasion, and metastases. Aberrant miR-106b expression is linked with breast cancer (BCa), prostate cancer (PCa),

E-mail address: ssagar.du@yahoo.com.

Peer review under responsibility of Chongqing Medical University.

lung cancer, gastric cancer (GC), colorectal cancer (CRC), hepatocellular carcinoma (HCC), and esophageal squamous cell carcinoma (ESCC). miR-106b and its seed family member like miR-106b-5p shows oncogenic properties and regulates different signaling pathways by targeting different genes involved in tumor initiation and progression and contributes to resistance against anti-cancer therapies.^{6,7} As its oncogenic role, several potential targets, like PTEN/AKT, La related protein 4B (LARP4B), Calponin 1, Runt-related transcription factor 3 (RUNX3), Disabled homolog 2 (DAB2), Deleted in liver cancer-1 (DLC-1), FOG2, RE-1 Silencing Transcription Factor (REST-1), Fucosyltransferase 6 (FUT6), Arm protein lost in epithelial cancers, on chromosome X (ALEX1) and BTG anti-proliferation factor 3 (BTG3) have been confirmed for their involvement in cancer initiation and progression through multiple pathways.

Moreover, miR-106b overexpression is associated with aggressive clinicopathological features. Experimental findings have shown that miR-106b could be an attractive and potential biomarker in different types of cancers for detection and prognosis assessment of the tumor. Clinical findings have reported that specific alteration in tumors in miRNAs plasma/serum of cancer patients has exhibited the plasma circulating miRNAs to be the new noninvasive biomarkers in various cancer patients.⁸ In the past decade, the importance of microRNAs (miRNAs) in cancer therapeutics have been progressively acknowledged. Particularly, miRNAs had been lately proven to reverse the resistance to chemotherapy drugs such as Doxorubicin, Cisplatin, and 5-fluorouracil.⁹

This review provides a brief overview of miR106b-mediated signaling pathways in different cancer types and focuses on the expression and targets of miR106 family members in cancer tissue or cells. The article further explores the possibilities of exploring the roles of this miRNA for diagnostic and prognostic purposes in cancer therapeutics.

miRNA biogenesis: mechanism and important players

The synthesis of miRNA is a multi-step process in which primary (pri-miRNA) is transcribed from the Minichromosome Maintenance Complex Component 7 (*MCM7*) gene by RNA polymerase.^{10–12} In some cases, miRNAs are transcribed as one lengthy transcript known as clusters, which may have similar seed regions. In such cases, these are considered as a family.¹³ Further, miRNA biogenesis includes cleavage events, one in the nucleus and another in the cytoplasm performed by RNase III family members, Drosha in the nucleus, and Dicer in the cytoplasm.¹⁴ The processing and cleavage of pri-miRNA are performed by a complex of Drosha/DGCR8 (*DiGeorge syndrome critical region 8*) to release the intermediate pre-miRNA. It binds to Exp5/Ran-GTP complex and gets exported to the cytoplasm. The processing of pri-miRNA depends on the structural characteristics of distinct pri-miRNA sequences.¹⁵ Next, pre-miRNA is digested by Dicer into a mature duplex (miRNA/miRNA*), where miRNA* represents a passenger strand.

Under normal circumstances, one strand from the duplex is recruited into the miRNA-induced silencing complex (miRISC) and functions as a guide for miRNA to recognize the mRNA targets and guide for translational repression, deadenylation, and degradation while another passenger strand is degraded. During this process, proteins like Argonaute and glycine–tryptophan protein 182 kDa (GW182) interact with miRNAs. The resultant RISC complex binds to the mRNA's 3'untranslated regions and plays a crucial role in miRNA-mediated mRNA degradation and translational repression^{16,17} (Fig. 1). The miRNA mediated gene regulation depends on the extent of the complementary pairing of miRNA and sequence of target mRNA. The partial homology between miRNA:mRNA target sequences lead to translation repression; however, a perfect homology may result in mRNA degradation.^{17,18}

miR-106b family members

The miR-106b family members -miR-106a/b, 17-5p, 20a/b, and 93, are overexpressed in multiple cancer and involves in cell cycle regulation and functioning of the checkpoint and promote cell cycle progression.¹⁹ miR-106b with miR-93 and -25 forms a cluster of miR-106b-25^{20–22} (Fig. 2). miR-106b-25 cluster is situated in the intron region of *MCM7* gene,²³ which is important for DNA replication initiation.²⁴ miR-106b-25 cluster has been reported to be overexpressed in multiple cancers, including HCC,²² esophageal adenocarcinoma,²⁵ BCa,²⁶ and prostate cancer.^{22,23,27} This cluster has been demonstrated as pro-oncogenic and shown to play a role in proliferation, anti-apoptotic, and cell cycle progression by regulating TGF- β .²⁸ Similar to miR-106b, miR-106b-5p transcribed from the cluster of miR-106b-25 is also reported as oncogene in BCa,^{29–31} GC,³² HCC.^{7,33,34} miR-106-5p function like other miRNAs by complementary pairing to specific mRNAs to degrade/suppress translation of targeted mRNA -promoting or -suppressing genes. Several targets of miR-106-5p have been reported in human cancer and play a major role in proliferation, migration, invasion, and cell cycle.³⁵

Regulation of miR-106b

Studies demonstrated that the miR-106b cluster is situated in the *MCM7* gene.²³ Aberrant expression of *MCM7* is seen in several human cancer³⁶ and correlated with the miR-106b cluster expression.³⁷ The transcription factor E2F has been shown to regulate the miR-106b cluster in several cancers. It has been reported that E2Fs have several transcriptional sites in *MCM7* promotor and directly regulate a few miRNAs cluster and in turn control E2F expression.^{38,39} Thangavel et al reported that *MCM7* is mainly regulated by Retinoblastoma (RB) through binding of the proximal E2F element in the promoter in breast cancer model⁴⁰ and suggested the miR-106b cluster repression through the RB/E2F pathway. However, the *MCM7* promoter also possesses an E-box binding site for the MYC.³⁹ Another study also showed that MYC oncogene regulates both the miR-106b-25 cluster and *MCM7* via E2Fs. The family of MYC oncogene, including MYC, MYCL, and MYCN, play major functions in tumor development and progression and therapeutic




Figure 1 A schematic of miRNAs biogenesis. miRNA biogenesis begins in the nucleus, where genes are transcribed to a primary miRNA (pri-miRNA) by RNA polymerase II. The pri-miRNA is processed and cleaved by Drosha/DGCR8 complex to release the intermediate pre-miRNA. It binds to Exp5/Ran-GTP complex and gets exported to the cytoplasm. Next, pre-miRNA is digested by Dicer into a mature duplex (miRNA/miRNA*), where miRNA* represents as a passenger strand. One strand of this duplex is loaded into the miRNA-induced silencing complex (miRISC) and functions as a guide for miRNA to recognize the mRNA targets and induce their translational repression deadenylation and degradation while another passenger strand is degraded. During this process, proteins like Argonaute and glycine–tryptophan protein 182 kDa (GW182) interacts with miRNAs. The resultant RISC complex binds to the mRNA’s 3’untranslated regions and plays a crucial role in miRNA-mediated mRNA degradation and translational repression.




Figure 2 A schematic illustration of the miR-106b-25 clusters. The genomic structure of the miR-106b-25 cluster on chromosome 7. Sequences of the miR-106b family members and conserved seed sequences among the family members are highlighted with blue color.

resistance.⁴¹ MYC activates transcription factors E2F1 and E2F3, which leads to increase MCM7 and miR-106b-25 clusters. In turn, E2F expression is downregulated by miR-106b and miR-93 through E2F1 transcript through negative feedback loop^{42,43} (Fig. 3).

MCM7 is also directly regulated by MYC family member-MYCN.⁴⁴ MYCN role has been reported in prostate neuroendocrine differentiation and frequently amplified in neuroblastoma.^{45,46} This report suggests that MYCN activates the MCM7 promoter through direct interaction.⁴⁴ Recently, bromodomain-containing protein 4 (BRD4) has been identified in the regulation of miR-106-5p transcription through

E2F³². In this study, BRD4 was found to be recruited to the promoter of miR-106b-5p with the help of E2F and aided miR-106b-5p transcription. In addition, BRD4 negatively regulates c-MYC stability and is involved in the cell cycle, inflammatory cytokines, and cancer development.^{32,47} Gong et al demonstrated the role of the transforming growth factor-beta (TGF-β) in miRNA regulation. The dual nature of TGF-β has been reported in many cancers, either as a tumor suppressor or tumor promoter. TGF-β upregulates transcription of miR-106b, independent of the promoter of the MCM7 gene by c-jun activation.⁴⁸ Hippo signaling pathway co-activator YAP/TAZ is known to




Figure 3 Oncogenic miR-106b regulation and its target. miR-106b-25 cluster is located in the intron region of the *MCM7* gene. miR-106b family is mainly regulated through E2Fs which have several transcriptional sites in *MCM7* promotor and directly regulate miR-106b family and in turn control E2F expression. MYC, MYCN, BRD4, and RB upregulate the expression of E2Fs. YAP/TAZ and TGF- β directly regulate the expression of miR-106 family, independent of *MCM7*. The miR-106b family targets multiple genes involved in different biological functions like proliferation, apoptosis, cell cycle, migration, invasion, and metastases.

regulate several miRNAs including miR-106b-25 cluster.^{49,50} In lung cancer, YAP/TAZ acts as an oncogene and regulates miR-106b-25 through *MCM7* gene and its hosted miRs transcription and regulates promote cell proliferation.⁴⁹

Breast cancer

BCa is by far the most common invasive cancer in females and the leading cause of cancer mortality worldwide.⁵¹ An estimated 276,480 new cases and 42,170 new deaths will be reported in the United States (US) in 2020.⁵² Breast cancer management employed various approaches like locoregional surgery and radiation therapy and systemic therapy.⁵³ However, some proportion of patients with early-stage of breast cancer develops recurrence in a few years.^{54,55} This explains a need for further improvement in the outcome and the problem of therapeutic resistance in breast cancer patients. Studies have found that miR-106b expression is elevated in breast tumor than non-tumor

tissues and associated with breast cancer progression. The author showed that miR-106b regulates several cellular processes in breast cancer *in vitro* and tumor growth *in vivo* and found that high miR-106b lowered the PTEN level in breast cancer and suggested that PTEN is immediate target of miR-106b. In addition, targeting AKT inhibits the ability of proliferation, migration and invasion ability in breast cancer cells. This suggest that miR-106b regulate cell progression through PTEN suppression in breast cancer. Moreover, upregulation of miR-106b promoted tumor growth in xenograft mouse model and *vice versa*. These results conclude that miR-106b promoted breast cancer progression and function as an oncogene.⁵⁶

Another study showed that miR-106b links with poor survival in invasive breast cancer patients. Zheng et al reported that miR-106b level is 2.4-fold higher in BCa tissue as compared to benign tumors and correlated with more advanced disease and bad prognosis in BCa patients. These results demonstrated that miR-106b role in tumor metastasis regulation and might be a prognostic marker for

recurrence in breast cancer.⁵⁵ miR-106b seed family member -miR-106-5p is found to be increased in breast cancer cells and activates Rho/ROCK1 pathway by targeting Calponin-1 in breast cancer cells. The author showed that inhibition of miR-106-5p led to a significantly enhanced CNN1 protein level or *vice versa* in breast cancer cells. Overexpression of CNN1 results in decreased cell proliferation, cell migration, and cell adhesion but accelerates the apoptosis rate in breast cancer cells. Forced overexpression of CNN1 led to a decrease in the mRNA and protein level of ROCK1 and protein level of Rho. Targeting the Rho/ROCK1 pathway relieves the effect of CNN1 on BCa cell proliferation and invasion. This suggests that miR-106-5p could lead to breast cancer progression.²⁹

In addition, the miR-106b expression is inversely proportional to the Fucosyltransferase 6 (FUT6), fucosyltransferase (FUT) family member leading to advanced cancer stages and poor prognosis in BCa. This study suggests that miR-106b inhibition increased the FUT6 expression and resulted in a decrease in cell proliferation, invasion, and migration.⁵⁷ The miR-106b-25 cluster activates the NOTCH1 signaling pathway and decreases E3 ubiquitin ligase expression- NEDD4L resulting in breast cancer initiation.²⁶ Interestingly, the authors showed that the miR-106b-5p expression could predict the progression and recurrence in BCa *in situ* via the TGF- β pathway.³⁰

Prostate cancer

PCa is noncutaneous cancer in men worldwide.⁵⁸ An estimated 191,930 new cases and 33,330 new deaths will be reported in the US in 2020.⁵² Despite the different therapies, advanced prostate cancer often develops therapeutic resistance to traditional therapies. Thus, a novel approach is needed to improve targeted therapies that target additional pathways in the growth and progression of prostate cancer. miRNAs have gained significant attention to prostate cancer progression.^{59,60} For example, miR-106b contributes to tumor viability and migration and acts as an oncogene in PCa. Similar to breast cancer, miR-106b is overexpressed and inversely proportional to PTEN in prostate cancer.²⁷ In prostate cancer, miR-106b expression is linked with tumor development and disease recurrence by targeting Caspase-7 and focal adhesion.⁶¹ Also, miR-106b has been shown to cause cell cycle arrest when p21 is downregulated in response to radiation therapy in PCa cells. Here, the authors demonstrated that expression of miR-106b increased three-fold after radiation and suppressed radiation-induced p21 activation in LNCaP prostate cancer cells. This study strongly supports that miR-106b directly targets and inhibits p21 and regulates G2/M cell cycle arrest.⁶²

Another study demonstrated that miR-106b possesses an anti-apoptotic role in PCa by targeting p21 and E2F1 and leads to inhibition of caspase activity.⁶³ Yin et al showed that miR-106b is overexpressed in prostate cancer tissue compared to paracancerous tissue. *LARP4B* (La related protein 4B), a LARP family member, is reported as a target gene and act as a tumor suppressor in PCa. *LARP4B* expression is shown to be lower and inversely proportional to miR-106b in PCa. miR-106b overexpression considerably

suppresses the cell viability, invasion, and migration of PCa cells by regulating the Ki67 expression (Proliferation marker), Matrix Metalloproteinase 2 (MMP-2), Cluster of Differentiation 44 (CD44), Smad2 in prostate cancer cells.⁶⁴ In addition, Liang et al also reported that miR-106b-25 expression is increased in LNCaP and PC-3 prostate cancer cells in hypoxia conditions and suppresses REST expression (RE-1 Silencing Transcription Factor) in this condition. This suggests the inverse relationship between miR-106b and REST in prostate cancer cells. Moreover, they showed that REST is downregulated and induction of at least one member of the miR-106b-25 cluster in high Gleason score (≥ 8) Prostate tumor analysis.⁶⁵

In the meta-analysis of human prostate biomarkers, miR-106b overexpression is significantly correlated with poor recurrence-free survival and suggested as a biomarker for diagnosing and predicting prostate cancer.^{66,67} Recently, a barely described RNA-binding protein (RBMS1) has been identified as a regulatory target of miR-106b. RBMS1 is shown to interact with miR-106b directly and is inversely proportional to miR-106b expression. Higher expression of miR-106b leads to loss of RBMS1 and influenced the cell growth, colony formation, and gap closing in prostate cancer tissue/cells.⁶⁸

Hepatocellular carcinoma

HCC is one of the aggressive types of liver cancer worldwide. HCC treatment selection depends on various aspects, like tumor characteristics, age, liver dysfunction, severity, and local expertise. Despite the recent advances in therapeutics, cancer-related death is continuing to rise worldwide.⁶⁹ Thus, identifying a new treatment approach and understanding the molecular mechanism involved in the HCC progression is needed. Several findings have shown that miR-106b and its member participates in HCC and associated with poor prognosis.⁷⁰⁻⁷² Yen et al showed that miR-106b expression is increased and linked with tumor progression and poor survival and increases recurrence rates in HBV-associated HCC patients. Moreover, patients with the well-differentiated HCC significantly showed lower expression of miR-106 compare to moderate and poor differentiation HCC patients, suggesting that miR-106b contributes to differentiation and poor survival with high recurrence chance.^{72,73}

Interestingly, overexpression of miR-106b upregulates the cell cycle regulator-Cyclin D1 through adenomatous polyposis coli (APC) downregulation to control the entry into the G1/S transitional phase.⁷⁴ Smad-7 is also identified as a miR-106b target and regulates the expression of TGF- β 1 and p-SMAD3 and results in miR-106b promoting the cell growth, migration, and EMT process through suppression of epithelial cell marker E-cadherin.⁷⁵ RhoGTPases are also identified as an indirect target of miR-106b and control stress fiber migration and cell migration.⁷⁶ Xu et al proposed that targeting miR-106b helps to overcome the tolerance of TNF-related apoptosis-inducing ligand (TRAIL) in HCC. They reported that miR-106b inhibitors increase Death receptor 4 (DR4) expression by direct targeting and augments TRAIL and therefore lower the acquired resistance to TRAIL.⁷⁷ POK transcription factor family 15

member - Zinc finger and BTB domain-containing protein 7A (Zbtb7a) is known to be the target of miR-106b, which plays a key role in tumorigenesis. miR-106b suppresses Zbtb7a translation and inhibits apoptosis.⁷⁸

Seed family member, miR-106-5p expression was found to be increased in HCC and shown to exhibit cancer stem cell properties and regulate cell migration by PTEN via PI3K/AKT pathway. Like other cancer types, miR-106-5p downregulates the PTEN expression in HCC and plays an important role in metastases in HCC cells, indicating the inverse relationship between miR-106b-5p level and PTEN. In addition, patients with a low level of PTEN are linked with more aggressive characteristics and more prolonged survival in patients with high PTEN expression.³³ In another study, HCC clinical showed that the higher expression of miR-106b-5p linked with bad prognosis and advanced TNM stages in patients. Functionally, miR-106b-5p accelerates the proliferation rate and invasion ability in HCC cells, and inhibition of miR-106b-5p by inhibitor lowers the cell proliferation rate and invasion ability. Runt-related transcription factor 3 (RUNX3), tumor suppressor, has been shown to direct miR-106-5p in HCC cells and functionally involved in the TGF- β mediated signaling pathway.^{79,80}

miR-106b overexpression enhances the cell viability and cell invasion by RUNX3, and targeting RUNX3 might reverse this phenotype.⁸¹ Yu et al also demonstrated overexpression of miR-106b-5p in HCC tissue associated with the patient's aggressive clinicopathological features. In addition, miR-106-5p overexpression promotes cell proliferation and invasion through FOG2 regulation. Furthermore, overexpression of miR-106b-5p decreases FOG2 (transcriptional regulator friend of Gata 2) expression and enhances cell proliferation and invasion of the HCC cells.⁷ miR-106b has been shown to target another tumor suppressor protein, disabled homolog 2 (DAB2). Expression of DAB2 is detected in HCC and plays a role in proliferation and migration. DAB2 is a negatively correlated and miR-106b direct target. Overexpression of miR-106b inhibits DAB2 and promotes the cell proliferation and ability of migration in HCC cells and suggested that miR-106b promote the HCC progression by DAB2 downregulation.⁸² Epidemiological studies have shown the role of hepatitis B "e" antigen (HBeAg) in the induction of miR-106b and enhance cell growth by targeting the retinoblastoma gene.⁸³ Circulating miR-106b in serum identified as a prognostic biomarker for HCC patients^{84,85} and in patients treated with Transcatheter arterial chemoembolization.⁸⁶

Lung cancer

Lung Cancer is a malignant tumor and one of the leading reasons for cancer mortality globally. An estimated 228,820 new cases and 135,720 deaths from lung cancer will be reported in the US in 2020.⁵² Despite the great improvements in lung cancer therapeutics, the prognosis remains poor. Most patients are diagnosed at stages III or IV before therapy, and the 5-year survival rate is less than 18%. Thus, there is a need to develop a detailed understating of tumor development from different aspects.^{87,88} Reports suggest that miR-106b is increased in lung cancer patients and cell lines and enhances invasion and migration in lung cancer

cells. Similarly, miR-106b directly regulated PTEN mRNA and protein expression in lung cancer cells and increase cell migration and invasion ability. In addition, they showed that the miR-106b serum exosomal level is positively correlated with TNM stages and lymph node metastasis.⁸⁹

Other studies showed that miR-106b -5p targets BTG anti-proliferation factor 3 (BTG3) in lung cancer cells. miR-106b-5p involves in tumor formation by suppressing the BTG3. Expression of BTG3 is reported to suppress proliferation, the progression of the cell cycle, and metastasis. The authors showed that miR-106b-5p is highly expressed in non-small-cell lung cancer (NSCLC) cells, promotes proliferation and caspase-mediated cell death inhibition by negatively regulating BTG3 through direct interaction and binding to BTG3-3'UTR.⁹⁰ In addition to regulating cancer initiation and progression, miR-106-5p is involved in the resistance to chemotherapy agents like cisplatin in NSCLC. This study demonstrated that miR-106b-5p directly targets polycystic kidney disease-2 (PKD2) through direct interaction in cisplatin-resistant NSCLC cells. This indicates that miR-106b-5p-mediated regulatory mechanism of cisplatin chemosensitivity NSCLC.⁹¹

The miR-106b cluster regulates the β -TRCP2 expression by directly targeting the β -TRCP2 transcript, augments cell invasion, and migration potential through Snail upregulation, which is known as β -TRCP2.⁹² Transforming Growth Factor- β (TGF- β) Receptor 2 (TGFBR2) shows tumor suppressor activity and is found to be downregulated in cancers, including NSCLC. The oncogenic miR-106b-25 cluster involves in TGFBR2 regulation in NSCLC and post-transcriptionally repress TGFBR2 level.⁵⁰

Gastric cancer

GC is the leading cause of cancer mortality globally with a poor prognosis.^{93,94} Treatment for gastric cancer includes surgery, radiation, chemotherapy, and neoadjuvant therapy, which shows improvement, but patients develop advanced stage due to low screening at early diagnosis.^{94,95} Due to poor prognosis, development of resistance to conventional therapy, late diagnosis, and search for biomarker remains a major obstacle in gastric cancer.^{9,96} Several studies reported that miRNAs dysregulation was found to be involved in gastric cancer progression.⁹⁷ The class of miRNAs has been identified as a prospective biomarker in GC early detection.⁹⁸ Among these miRNAs, miR-106b significantly upregulated in the gastric cancer^{8,99} and genetic association with gastric cancer.¹⁰⁰ In addition, large scale analysis, miR-106b plasma concentration is significantly higher in GC patients as compared to control.¹⁰¹ Studies have shown that miR-106b alters the feature of cancer stem cell (CSC) by direct regulation of SMAD7 and subsequently inhibits TGF- β in CD44-positive GC cells.¹⁰² Yang et al also demonstrated that miR-106b expression is higher in cancer-associated fibroblasts (CAFs) from GC and linked with poor prognosis, promote cell migration and invasion through PTEN.¹⁰³

Recently, miR-106b role in metastasis of early GC has been reported and found that miR-106b expression is more in the metastasis tissues compared with non-metastasis tissues. In this study, miR-106b directly targets a novel

tumor suppressor gene, *ALEX1*, which is known to suppress gastric cancer metastasis and inhibit apoptosis through the JAK/STAT pathway both *in vitro* and *in vivo*.¹⁰⁴ Studies have shown that miR-106b-25 cluster, especially miR-106b, is upregulated in gastric tumor subsets and correlated with lymph node metastases, distant metastasis, and TNM stage^{43,105,106} and regulated by E2F1; in turn, miR-106b and miR-93 downregulates the E2F1 expression through negative feedback loop.^{42,43} Furthermore, miR-106b upregulation impairs the TGF- β pathway and regulates CDKN1A and BIM and suggested that the miR-106b-25 cluster is involved in TGF- β resistance development.⁴³ In addition to TGF- β resistance, miR-106-5p has been reported to be involved in resistance *via* TNF-related apoptosis-inducing ligand (TRAIL), known to promote apoptosis in cancer cells.¹⁰⁷

Esophageal squamous cell carcinoma

In ESCC, miR-106b works as an oncogene and promotes proliferation, invasion, and metastases. Recently, Zhang et al showed that miR-106b expression enhanced in ESCC tissues. Functionally, they showed that targeting miR-106 with inhibitor significantly decreased wound healing ability as compared to control groups in EC-1 cells. Also, miR-106 inhibition could inhibit the induction of EMT in ESCC cells. Furthermore, they demonstrated that miR-106b directly targets PTEN and involves in invasion, metastases, and proliferation. Downregulation of PTEN in miR-106b treated cells rescued the EMT induction in ESCC cells and suggested that miR-106b mediated EMT depends on PTEN expression.¹⁰⁸ In recent years, increased miR-106b expression is involved in lymph node (LN) metastases and promotes cell invasion and migration through epithelial–mesenchymal transition (EMT) *via* SMAD-7 downregulation.¹⁰⁹

Similarly, another study proved that miR-106b is linked with the clinicopathological features of ESCC. This study showed that miR-106b overexpression is associated with smoking, TNM classification, and LN metastasis and smoking. Also, low miR-106b expression in patients showed a higher survival rate compare to patients with increased expression.¹¹⁰

Colorectal cancer

CRC is the most lethal cancer and the third leading cause of cancer mortality globally.^{111,112} Overall, colorectal associated death rates have dropped over the years due to early detection because this CRC is curable in its early stage, but incidence rates still remain high.^{113,114} Surgery followed by chemotherapy is the first line of treatment in Colorectal cancer, but the prognosis is never favorable.¹¹¹ Targeted therapy for advanced-stage tumors has shown an increased survival rate,¹¹¹ and new drugs have shown less success and develop resistance.^{111,114,115} Several studies showed the role of aberrant expression of miRNAs in CRC initiation, progression, and metastases.^{113,116} Zhang et al reported that miR-106b is significantly overexpressed in tissues of metastatic CRC and linked with advanced clinical CRC stage and LN metastasis. Metastatic suppressor gene *DLC-1* has been recognized as miR-106b direct target and inversely

Table 1 miR-106b targets and role in cancer types.

Cancer types	Target genes	Biological role	Reference
Breast Cancer	<i>PTEN</i>	Tumor growth	56
	<i>Calponin-1</i>	Cell proliferation and invasion	29
	<i>TGF-β</i>	Proliferation	30
	<i>FUT6</i>	Proliferation, invasion, and migration	57
Prostate Cancer	<i>PTEN</i>	Cell viability and migration	27
	<i>Caspase-7</i>	Tumor development and disease recurrence	61
	<i>P21, E2F1</i>	Cell cycle arrest, anti-apoptotic	62, 63
	<i>LARP4B</i>	Tumor viability and metastases	64
	<i>REST-1</i>	Neuronal markers	65
	<i>RBMS1</i>	Growth, colony formation & gap closing	68
Hepatocellular Carcinoma	<i>PTEN</i>	Aggressiveness and disease-free survival	33
	<i>Cyclin-D1</i>	Cell cycle	74
	<i>SMAD7</i>	Cell growth, migration and EMT	75
	<i>Rho/GTPases</i>	Migration	76
	<i>TRAIL</i>	Proliferation, apoptosis, and resistance	77
	<i>Zbtb7a</i>	Apoptosis	78
	<i>RUNX3</i>	Proliferation, invasion, migration and angiogenesis	79–81
	<i>FOG2</i>	Proliferation and invasion	7
Lung Cancer	<i>DAB2</i>	Proliferation and migration	82
	<i>PTEN</i>	Migration and invasion	89
	<i>BTG3</i>	Proliferation and anti-apoptosis	90
	<i>PKD2</i>	Resistance to chemotherapy	91
	<i>β-TRCP2</i>	Invasion and migration	92
Gastric Cancer	<i>TGFBR2</i>	Cell proliferation, apoptosis	50
	<i>E2F1</i>	Apoptosis, Metastases, TNM stages	42,43
	<i>CDKN1A & BIM</i>	Cell cycle arrest and Apoptosis	43
	<i>SMAD7</i>	Cancer stem cell characteristics	102
	<i>PTEN</i>	Proliferation, invasion	103

(continued on next page)

Table 1 (continued)

Cancer types	Target genes	Biological role	Reference
ESCC		and migration	
	ALEX1	Metastasis, apoptosis	104
	TRAIL	Apoptosis	107
	PTEN	Proliferation, invasion, migration, EMT	108
CRC	SMAD7	Invasion, migration and metastasis	109
	DLC-1	Cell migration and invasion	117
	PTEN, P21	Cell radioresistance	118

correlated in CRC tissues. Overexpression of miR-106b enhances cell migration and invasion via DLC-1¹¹⁷. In addition, miR-106b involved in cell radioresistance via its direct target PTEN and p21 in colorectal cancer in *in vivo* & *in vitro* studies.¹¹⁸ Recently, a meta-analysis study suggested that the miR-106 family is participating in the CRC initiation, development, and progression and a promising therapeutic biomarker for CRC in meta-analysis.¹¹⁹

Conclusions

In recent years, miR-106b and its family members have emerged as putative oncogenes that can target multiple genes and promote tumor progression via different cancer signaling pathways (Table 1). This review discussed the evidence of miR-106b family members' role in cancer and contributed to resistance against anti-cancer therapies. Despite the advances in therapy, cancer cells develop resistance to therapies such as radiation therapy, chemotherapy, and targeted therapies by different mechanisms, including genetics or epigenetic changes in the cancer cell. Recently, miRNAs have become known as a prospective biomarker in different cancer types and their therapeutic potential. Expression of miRNAs has shown to increase in different cancer types and involve in cancer progression. In the current era of conventional therapy, miRNA-based therapies show a novel approach in cancer management. miRNA-based therapy like miRNA mimics and antagomir has been developed to modulate the tumor microenvironment and inhibits tumor progression.

Moreover, miRNA-based therapies have progressed in clinical trials and will provide more understanding of this new cancer therapeutic approach. There are some challenges in this approach, like 1) effectiveness of miRNA-based therapy depends on the precise, efficient and suitable delivery of miRNA and 2) comprehensive understanding of the biological roles of circulating miRNAs are a potential barrier. Despite all the challenges, the miRNA targeted approach holds strong potential in cancer therapeutics. The conventional therapies are partially effective and/or develop resistance later; combinatorial approaches with miRNA-based therapy could open new cancer therapeutics strategies. miR-106b and other family members

could be attractive and potential biomarkers in different cancer types for detection, diagnosis, and prognosis assessment in the tumor. In summary, targeting miR-106 with novel approaches could become potential strategies in patients who develop resistance to conventional therapies to overcome the resistance and bring new hope to cancer patients.

Conflict of interests

The author declares no conflict of interest.

References

- Samanta S, Balasubramanian S, Rajasingh S, et al. MicroRNA: a new therapeutic strategy for cardiovascular diseases. *Trends Cardiovasc Med.* 2016;26(5):407–419.
- Markou A, Zavridou M, Lianidou ES. miRNA-21 as a novel therapeutic target in lung cancer. *Lung Cancer (Auckl).* 2016; 7:19–27.
- Peng Y, Croce CM. The role of MicroRNAs in human cancer. *Signal Transduct Target Ther.* 2016;1:e15004.
- Ha M, Kim VN. Regulation of microRNA biogenesis. *Nat Rev Mol Cell Biol.* 2014;15(8):509–524.
- Filipow S, Laczmanski L. Blood circulating miRNAs as cancer biomarkers for diagnosis and surgical treatment response. *Front Genet.* 2019;10:e169.
- Jiao D, Yan Y, Shui S, et al. miR-106b regulates the 5-fluorouracil resistance by targeting Zbtb7a in cholangiocarcinoma. *Oncotarget.* 2017;8(32):52913–52922.
- Yu LX, Zhang BL, Yang MY, et al. MicroRNA-106b-5p promotes hepatocellular carcinoma development via modulating FOG2. *Onco Targets Ther.* 2019;12:5639–5647.
- Gorur A, Balci Fidanci S, Dogruer Unal N, et al. Determination of plasma microRNA for early detection of gastric cancer. *Mol Biol Rep.* 2013;40(3):2091–2096.
- Zhao X, Hu GF, Shi YF, Xu W. Research progress in microRNA-based therapy for gastric cancer. *Onco Targets Ther.* 2019;12: 11393–11411.
- Macfarlane LA, Murphy PR. MicroRNA: biogenesis, function and role in cancer. *Curr Genom.* 2010;11(7): 537–561.
- Bartel DP. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell.* 2004;116(2):281–297.
- Lee Y, Kim M, Han J, et al. MicroRNA genes are transcribed by RNA polymerase II. *EMBO J.* 2004;23(20):4051–4060.
- Tanzer A, Stadler PF. Molecular evolution of a microRNA cluster. *J Mol Biol.* 2004;339(2):327–335.
- Kim VN, Han J, Siomi MC. Biogenesis of small RNAs in animals. *Nat Rev Mol Cell Biol.* 2009;10(2):126–139.
- O'Carroll D, Schaefer A. General principals of miRNA biogenesis and regulation in the brain. *Neuro-psychopharmacology.* 2013;38(1):39–54.
- Treiber T, Treiber N, Meister G. Regulation of microRNA biogenesis and its crosstalk with other cellular pathways. *Nat Rev Mol Cell Biol.* 2019;20(1):5–20.
- O'Brien J, Hayder H, Zayed Y, Peng C. Overview of MicroRNA biogenesis, mechanisms of actions, and circulation. *Front Endocrinol (Lausanne).* 2018;9:e402.
- Fabian MR, Sonenberg N. The mechanics of miRNA-mediated gene silencing: a look under the hood of miRISC. *Nat Struct Mol Biol.* 2012;19(6):586–593.
- Ivanovska I, Ball AS, Diaz RL, et al. MicroRNAs in the miR-106b family regulate p21/CDKN1A and promote cell cycle progression. *Mol Cell Biol.* 2008;28(7):2167–2174.

20. Petrocca F, Vecchione A, Croce CM. Emerging role of miR-106b-25/miR-17-92 clusters in the control of transforming growth factor beta signaling. *Cancer Res.* 2008;68(20):8191–8194.
21. Mendell JT. miRiad roles for the miR-17-92 cluster in development and disease. *Cell.* 2008;133(2):217–222.
22. Li Y, Tan W, Neo TW, et al. Role of the miR-106b-25 microRNA cluster in hepatocellular carcinoma. *Cancer Sci.* 2009;100(7):1234–1242.
23. Khuu C, Utheim TP, Sehic A. The three paralogous MicroRNA clusters in development and disease, miR-17-92, miR-106a-363, and miR-106b-25. *Scientifica (Cairo).* 2016;2016:e1379643.
24. Mehlich D, Garbicz F, Wlodarski PK. The emerging roles of the polycistronic miR-106b approximately 25 cluster in cancer - a comprehensive review. *Biomed Pharmacother.* 2018;107:1183–1195.
25. Kan T, Sato F, Ito T, et al. The miR-106b-25 polycistron, activated by genomic amplification, functions as an oncogene by suppressing p21 and Bim. *Gastroenterology.* 2009;136(5):1689–1700.
26. Guarnieri AL, Towers CG, Drasin DJ, et al. The miR-106b-25 cluster mediates breast tumor initiation through activation of NOTCH1 via direct repression of NEDD4L. *Oncogene.* 2018;37(28):3879–3893.
27. Poliseno L, Salmena L, Riccardi L, et al. Identification of the miR-106b-25 microRNA cluster as a proto-oncogenic PTEN-targeting intron that cooperates with its host gene MCM7 in transformation. *Sci Signal.* 2010;3(117):ra29.
28. Li F, Liu J, Li S. MicorRNA 106b approximately 25 cluster and gastric cancer. *Surg Oncol.* 2013;22(2):e7–e10.
29. Wang Z, Li TE, Chen M, Pan JJ, Shen KW. miR-106b-5p contributes to the lung metastasis of breast cancer via targeting CNN1 and regulating Rho/ROCK1 pathway. *Aging (Albany NY).* 2020;12(2):1867–1887.
30. Lee J, Kim HE, Song YS, Cho EY, Lee A. miR-106b-5p and miR-17-5p could predict recurrence and progression in breast ductal carcinoma in situ based on the transforming growth factor-beta pathway. *Breast Cancer Res Treat.* 2019;176(1):119–130.
31. Banerjee S, Kalyani Yabalooru SR, Karunagaran D. Identification of mRNA and non-coding RNA hubs using network analysis in organ tropism regulated triple negative breast cancer metastasis. *Comput Biol Med.* 2020;127:e104076.
32. Dong X, Hu X, Chen J, Hu D, Chen LF. BRD4 regulates cellular senescence in gastric cancer cells via E2F/miR-106b/p21 axis. *Cell Death Dis.* 2018;9(2):230.
33. Shi DM, Bian XY, Qin CD, Wu WZ. miR-106b-5p promotes stem cell-like properties of hepatocellular carcinoma cells by targeting PTEN via PI3K/Akt pathway. *Onco Targets Ther.* 2018;11:571–585.
34. Zhu HR, Huang RZ, Yu XN, et al. Microarray expression profiling of microRNAs reveals potential biomarkers for hepatocellular carcinoma. *Tohoku J Exp Med.* 2018;245(2):89–98.
35. Yang C, Dou R, Yin T, Ding J. MiRNA-106b-5p in human cancers: diverse functions and promising biomarker. *Biomed Pharmacother.* 2020;127:e110211.
36. Simon NE, Schwacha A. The Mcm2-7 replicative helicase: a promising chemotherapeutic target. *Biomed Res Int.* 2014;2014:e549719.
37. Markey MP, Angus SP, Strobeck MW, et al. Unbiased analysis of RB-mediated transcriptional repression identifies novel targets and distinctions from E2F action. *Cancer Res.* 2002;62(22):6587–6597.
38. Woods K, Thomson JM, Hammond SM. Direct regulation of an oncogenic micro-RNA cluster by E2F transcription factors. *J Biol Chem.* 2007;282(4):2130–2134.
39. Suzuki S, Adachi A, Hiraiwa A, Ohashi M, Ishibashi M, Kiyono T. Cloning and characterization of human MCM7 promoter. *Gene.* 1998;216(1):85–91.
40. Thangavel C, Boopathi E, Ertel A, et al. Regulation of miR106b cluster through the RB pathway: mechanism and functional targets. *Cell Cycle.* 2013;12(1):98–111.
41. Han H, Jain AD, Truica MI, et al. Small-molecule MYC inhibitors suppress tumor growth and enhance immunotherapy. *Cancer Cell.* 2019;36(5):483–497.
42. Bueno MJ, Gomez de Cedron M, Laresgoiti U, Fernandez-Piqueras J, Zubiaga AM, Malumbres M. Multiple E2F-induced microRNAs prevent replicative stress in response to mitogenic signaling. *Mol Cell Biol.* 2010;30(12):2983–2995.
43. Petrocca F, Visone R, Onelli MR, et al. E2F1-regulated microRNAs impair TGFbeta-dependent cell-cycle arrest and apoptosis in gastric cancer. *Cancer Cell.* 2008;13(3):272–286.
44. Shohet JM, Hicks MJ, Plon SE, et al. Minichromosome maintenance protein MCM7 is a direct target of the MYCN transcription factor in neuroblastoma. *Cancer Res.* 2002;62(4):1123–1128.
45. Carneiro BA, Pamarthy S, Shah AN, et al. Anaplastic lymphoma kinase mutation (ALK F1174C) in small cell carcinoma of the prostate and molecular response to alectinib. *Clin Cancer Res.* 2018;24(12):2732–2739.
46. Huang M, Weiss WA. Neuroblastoma and MYCN. *Cold Spring Harb Perspect Med.* 2013;3(10):a014415.
47. Devaiah BN, Mu J, Akman B, et al. MYC protein stability is negatively regulated by BRD4. *Proc Natl Acad Sci U S A.* 2020;117(24):13457–13467.
48. Gong C, Qu S, Liu B, et al. MiR-106b expression determines the proliferation paradox of TGF-beta in breast cancer cells. *Oncogene.* 2015;34(1):84–93.
49. Lo Sardo F, Forcato M, Sacconi A, et al. MCM7 and its hosted miR-25, 93 and 106b cluster elicit YAP/TAZ oncogenic activity in lung cancer. *Carcinogenesis.* 2017;38(1):64–75.
50. Lo Sardo F, Pulito C, Sacconi A, et al. YAP/TAZ and EZH2 synergize to impair tumor suppressor activity of TGFBR2 in non-small cell lung cancer. *Cancer Lett.* 2020;500:51–63.
51. Ataollahi MR, Sharifi J, Paknahad MR, Paknahad A. Breast cancer and associated factors: a review. *J Med Life.* 2015;8(Spec Iss 4):6–11.
52. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2020. *CA Cancer J Clin.* 2020;70(1):7–30.
53. Harbeck N, Penault-Llorca F, Cortes J, et al. Breast cancer. *Nat Rev Dis Primers.* 2019;5(1):e66.
54. Bosco JL, Lash TL, Prout MN, et al. Breast cancer recurrence in older women five to ten years after diagnosis. *Cancer Epidemiol Biomarkers Prev.* 2009;18(11):2979–2983.
55. Zheng R, Pan L, Gao J, et al. Prognostic value of miR-106b expression in breast cancer patients. *J Surg Res.* 2015;195(1):158–165.
56. Li N, Miao Y, Shan Y, et al. MiR-106b and miR-93 regulate cell progression by suppression of PTEN via PI3K/Akt pathway in breast cancer. *Cell Death Dis.* 2017;8(5):2796.
57. Li N, Liu Y, Miao Y, Zhao L, Zhou H, Jia L. MicroRNA-106b targets FUT6 to promote cell migration, invasion, and proliferation in human breast cancer. *IUBMB Life.* 2016;68(9):764–775.
58. Sagar V, Vatapalli R, Lysy B, et al. EPHB4 inhibition activates ER stress to promote immunogenic cell death of prostate cancer cells. *Cell Death Dis.* 2019;10(11):801.
59. Malla B, Aebersold DM, Dal Pra A. Protocol for serum exosomal miRNAs analysis in prostate cancer patients treated with radiotherapy. *J Transl Med.* 2018;16(1):e223.
60. Xiaoli Z, Yawei W, Lianna L, Haifeng L, Hui Z. Screening of target genes and regulatory function of miRNAs as prognostic

- indicators for prostate cancer. *Med Sci Monit.* 2015;21:3748–3759.
61. Hudson RS, Yi M, Esposito D, et al. MicroRNA-106b-25 cluster expression is associated with early disease recurrence and targets caspase-7 and focal adhesion in human prostate cancer. *Oncogene.* 2013;32(35):4139–4147.
 62. Li B, Shi XB, Nori D, et al. Down-regulation of microRNA 106b is involved in p21-mediated cell cycle arrest in response to radiation in prostate cancer cells. *Prostate.* 2011;71(6):567–574.
 63. Ambs S, Prueitt RL, Yi M, et al. Genomic profiling of microRNA and messenger RNA reveals deregulated microRNA expression in prostate cancer. *Cancer Res.* 2008;68(15):6162–6170.
 64. Yin W, Chen J, Wang G, Zhang D. MicroRNA106b functions as an oncogene and regulates tumor viability and metastasis by targeting LARP4B in prostate cancer. *Mol Med Rep.* 2019;20(2):951–958.
 65. Liang H, Studach L, Hullinger RL, Xie J, Andrisani OM. Down-regulation of RE-1 silencing transcription factor (REST) in advanced prostate cancer by hypoxia-induced miR-106b-25. *Exp Cell Res.* 2014;320(2):188–199.
 66. Song CJ, Chen H, Chen LZ, Ru GM, Guo JJ, Ding QN. The potential of microRNAs as human prostate cancer biomarkers: a meta-analysis of related studies. *J Cell Biochem.* 2018;119(3):2763–2786.
 67. Jin W, Fei X, Wang X, Chen F, Song Y. Circulating miRNAs as biomarkers for prostate cancer diagnosis in subjects with benign prostatic hyperplasia. *J Immunol Res.* 2020;2020:e5873056.
 68. Dankert JT, Wiesehofer M, Wach S, Czyrnik ED, Wennemuth G. Loss of RBMS1 as a regulatory target of miR-106b influences cell growth, gap closing and colony forming in prostate carcinoma. *Sci Rep.* 2020;10(1):18022.
 69. Yang JD, Hainaut P, Gores GJ, Amadou A, Plymoth A, Roberts LR. A global view of hepatocellular carcinoma: trends, risk, prevention and management. *Nat Rev Gastroenterol Hepatol.* 2019;16(10):589–604.
 70. Shen G, Jia H, Chen D, Zhang J. Effects of miR-106b expression on the proliferation of human hepatocellular carcinoma cells. *Zhonghua Zhong Liu Za Zhi.* 2014;36(7):489–495.
 71. Li BK, Huang PZ, Qiu JL, Liao YD, Hong J, Yuan YF. Upregulation of microRNA-106b is associated with poor prognosis in hepatocellular carcinoma. *Diagn Pathol.* 2014;9:e226.
 72. Zheng Q, Wei X, Rao J, Zhou C. Identification of key miRNAs in the progression of hepatocellular carcinoma using an integrated bioinformatics approach. *PeerJ.* 2020;8:e9000.
 73. Yen CS, Su ZR, Lee YP, Liu IT, Yen CJ. miR-106b promotes cancer progression in hepatitis B virus-associated hepatocellular carcinoma. *World J Gastroenterol.* 2016;22(22):5183–5192.
 74. Shen G, Jia H, Tai Q, Li Y, Chen D. miR-106b downregulates adenomatous polyposis coli and promotes cell proliferation in human hepatocellular carcinoma. *Carcinogenesis.* 2013;34(1):211–219.
 75. Li X, Zeng X. Shikonin suppresses progression and epithelial-mesenchymal transition in hepatocellular carcinoma (HCC) cells by modulating miR-106b/SMAD7/TGF-beta signaling pathway. *Cell Biol Int.* 2020;44(2):467–476.
 76. Yau WL, Lam CS, Ng L, et al. Over-expression of miR-106b promotes cell migration and metastasis in hepatocellular carcinoma by activating epithelial-mesenchymal transition process. *PLoS One.* 2013;8(3):e57882.
 77. Xu C, Shi L, Chen W, et al. MiR-106b inhibitors sensitize TRAIL-induced apoptosis in hepatocellular carcinoma through increase of death receptor 4. *Oncotarget.* 2017;8(26):41921–41931.
 78. Liang X, Zhao Q, Geng T, Luo S, He Q. MiR-106b regulates the apoptosis and tumorigenesis of hepatocellular carcinoma via targeting Zinc finger and BTB domain-containing protein 7A (Zbtb7A). *J Biochem Mol Toxicol.* 2018;32(8):e22169.
 79. Zhou WN, Du YF, Bai J, et al. RUNX3 plays a tumor suppressor role by inhibiting cell migration, invasion and angiogenesis in oral squamous cell carcinoma. *Oncol Rep.* 2017;38(4):2378–2386.
 80. Chen LF. Tumor suppressor function of RUNX3 in breast cancer. *J Cell Biochem.* 2012;113(5):1470–1477.
 81. Gu H, Gu S, Zhang X, et al. miR-106b-5p promotes aggressive progression of hepatocellular carcinoma via targeting RUNX3. *Cancer Med.* 2019;8(15):6756–6767.
 82. Sun C, Yao X, Jiang Q, Sun X. miR-106b targets DAB2 to promote hepatocellular carcinoma cell proliferation and metastasis. *Oncol Lett.* 2018;16(3):3063–3069.
 83. Samal J, Kandpal M, Vivekanandan P. HBeAg-induced miR-106b promotes cell growth by targeting the retinoblastoma gene. *Sci Rep.* 2017;7(1):14371.
 84. Shi BM, Lu W, Ji K, Wang YF, Xiao S, Wang XY. Study on the value of serum miR-106b for the early diagnosis of hepatocellular carcinoma. *World J Gastroenterol.* 2017;23(20):3713–3720.
 85. Yang DZ. Investigation on the estimation of bone mineral content in vivo by radiographic methods. *Zhonghua Fang She Xue Za Zhi.* 1985;19(3):149–152.
 86. Ali HEA, Emam AA, Zeeneldin AA, et al. Circulating miR-26a, miR-106b, miR-107 and miR-133b stratify hepatocellular carcinoma patients according to their response to transarterial chemoembolization. *Clin Biochem.* 2019;65:45–52.
 87. Lung cancer: despite advances, prevention is still best. *Lancet.* 2016;388(10044):e533.
 88. Hirono T, Jingushi K, Nagata T, et al. MicroRNA-130b functions as an oncomiRNA in non-small cell lung cancer by targeting tissue inhibitor of metalloproteinase-2. *Sci Rep.* 2019;9(1):6956.
 89. Sun S, Chen H, Xu C, et al. Exosomal miR-106b serves as a novel marker for lung cancer and promotes cancer metastasis via targeting PTEN. *Life Sci.* 2020;244:117297.
 90. Wei K, Pan C, Yao G, et al. MiR-106b-5p promotes proliferation and inhibits apoptosis by regulating BTG3 in non-small cell lung cancer. *Cell Physiol Biochem.* 2017;44(4):1545–1558.
 91. Yu S, Qin X, Chen T, Zhou L, Xu X, Feng J. MicroRNA-106b-5p regulates cisplatin chemosensitivity by targeting polycystic kidney disease-2 in non-small-cell lung cancer. *Anticancer Drugs.* 2017;28(8):852–860.
 92. Savita U, Karunakaran D. MicroRNA-106b-25 cluster targets beta-TRCP2, increases the expression of Snail and enhances cell migration and invasion in H1299 (non small cell lung cancer) cells. *Biochem Biophys Res Commun.* 2013;434(4):841–847.
 93. den Hoed CM, Kuipers EJ. Gastric cancer: how can we reduce the incidence of this disease? *Curr Gastroenterol Rep.* 2016;18(7):e34.
 94. Smyth EC, Nilsson M, Grabsch HI, van Grieken NC, Lordick F. Gastric cancer. *Lancet.* 2020;396(10251):635–648.
 95. Song Z, Wu Y, Yang J, Yang D, Fang X. Progress in the treatment of advanced gastric cancer. *Tumour Biol.* 2017;39(7):e1010428317714626.
 96. Shekari N, Javadian M, Ghasemi M, Baradaran B, Darabi M, Kazemi T. Synergistic beneficial effect of docosahexaenoic acid (DHA) and docetaxel on the expression level of Matrix metalloproteinase-2 (MMP-2) and MicroRNA-106b in gastric cancer. *J Gastrointest Cancer.* 2020;51(1):70–75.
 97. P LA, Ahadi A, Zare A, et al. Up-Regulation of miR-21, miR-25, miR-93, and miR-106b in Gastric Cancer. *Iran Biomed J.* 2018;22(6):367–373.
 98. Bhat SA, Majid S, Rehman MU. Scenario and future prospects of microRNAs in gastric cancer: a review. *Iran J Basic Med Sci.* 2019;22(4):345–352.

99. Bispo MA, Ramos P, Laranjeiro T, Cabral P, Palminha JM. Clinical and laboratory basis for the diagnosis and understanding of neonatal metabolic diseases. *Acta Med Port.* 1998;11(7):675–681.
100. Espinosa-Parrilla Y, Munoz X, Bonet C, et al. Genetic association of gastric cancer with miRNA clusters including the cancer-related genes MIR29, MIR25, MIR93 and MIR106: results from the EPIC-EURGAST study. *Int J Cancer.* 2014;135(9):2065–2076.
101. Tsujiura M, Ichikawa D, Komatsu S, et al. Circulating microRNAs in plasma of patients with gastric cancers. *Br J Cancer.* 2010;102(7):1174–1179.
102. Yu D, Shin HS, Lee YS, Lee YC. miR-106b modulates cancer stem cell characteristics through TGF-beta/Smad signaling in CD44-positive gastric cancer cells. *Lab Invest.* 2014;94(12):1370–1381.
103. Yang TS, Yang XH, Chen X, et al. MicroRNA-106b in cancer-associated fibroblasts from gastric cancer promotes cell migration and invasion by targeting PTEN. *FEBS Lett.* 2014;588(13):2162–2169.
104. Zhu Z, Yang Q, Zhang B, Wu W, Yuan F, Zhu Z. miR-106b promotes metastasis of early gastric cancer by targeting ALEX1 in vitro and in vivo. *Cell Physiol Biochem.* 2019;52(3):606–616.
105. Li F, Guo Y, Liu J, Zhang R. The significance of elevated plasma expression of microRNA 106b-25 clusters in gastric cancer. *PloS One.* 2017;12(5):e0178427.
106. Zhang R, Wang W, Li F, Zhang H, Liu J. MicroRNA-106b-25 expressions in tumor tissues and plasma of patients with gastric cancers. *Med Oncol.* 2014;31(10):e243.
107. Pereira J, Santos M, Delabio R, et al. Analysis of gene expression of miRNA-106b-5p and TRAIL in the apoptosis pathway in gastric cancer. *Genes (Basel).* 2020;11(4):e393.
108. Zhang J, Chen D, Liang S, et al. miR-106b promotes cell invasion and metastasis via PTEN mediated EMT in ESCC. *Oncol Lett.* 2018;15(4):4619–4626.
109. Dai F, Liu T, Zheng S, et al. MiR-106b promotes migration and invasion through enhancing EMT via downregulation of Smad 7 in Kazakh's esophageal squamous cell carcinoma. *Tumour Biol.* 2016;37(11):14595–14604.
110. Xiao CY, Guo ZJ, Cao L, Ding XW, Liu Q, Wang XC. Expression of mir-106b in esophageal squamous cell carcinoma. *Nan Fang Yi Ke Da Xue Xue Bao.* 2016;36(12):1667–1671.
111. Xie YH, Chen YX, Fang JY. Comprehensive review of targeted therapy for colorectal cancer. *Signal Transduct Target Ther.* 2020;5(1):e22.
112. Arnold M, Sierra MS, Laversanne M, Soerjomataram I, Jemal A, Bray F. Global patterns and trends in colorectal cancer incidence and mortality. *Gut.* 2017;66(4):683–691.
113. To KK, Tong CW, Wu M, Cho WC. MicroRNAs in the prognosis and therapy of colorectal cancer: from bench to bedside. *World J Gastroenterol.* 2018;24(27):2949–2973.
114. Strubberg AM, Madison BB. MicroRNAs in the etiology of colorectal cancer: pathways and clinical implications. *Dis Model Mech.* 2017;10(3):197–214.
115. Gong J, Cho M, Fakhri M. RAS and BRAF in metastatic colorectal cancer management. *J Gastrointest Oncol.* 2016;7(5):687–704.
116. Michael MZ, SMOC, van Holst Pellekaan NG, Young GP, James RJ. Reduced accumulation of specific microRNAs in colorectal neoplasia. *Mol Cancer Res.* 2003;1(12):882–891.
117. Zhang GJ, Li JS, Zhou H, Xiao HX, Li Y, Zhou T. MicroRNA-106b promotes colorectal cancer cell migration and invasion by directly targeting DLC1. *J Exp Clin Cancer Res.* 2015;34(1):e73.
118. Zheng L, Zhang Y, Liu Y, et al. MiR-106b induces cell radioresistance via the PTEN/PI3K/AKT pathways and p21 in colorectal cancer. *J Transl Med.* 2015;13:e252.
119. Peng Q, Shen Y, Zhao P, Cheng M, Zhu Y, Xu B. Biomarker roles identification of miR-106 family for predicting the risk and poor survival of colorectal cancer. *BMC Cancer.* 2020;20(1):e506.