



Two-proton radioactivity from excited states of proton-rich nuclei within Coulomb and Proximity Potential Model

De-Xing Zhu¹ · Yang-Yang Xu¹ · Li-Jia Chu² · Xi-Jun Wu³ · Biao He⁴ · Xiao-Hua Li^{1,5,6,7}

Received: 28 December 2022 / Revised: 8 June 2023 / Accepted: 25 June 2023 / Published online: 8 September 2023

© The Author(s), under exclusive licence to China Science Publishing & Media Ltd. (Science Press), Shanghai Institute of Applied Physics, the Chinese Academy of Sciences, Chinese Nuclear Society 2023

Abstract

In the present work, we extend the Coulomb and Proximity Potential Model (CPPM) to study two-proton (2p) radioactivity from excited states while the proximity potential is chosen as AW95 proposed by Aage Withner in 1995. Demonstration reveals that the theoretical results acquired by CPPM exhibit a high level of consistency with prior theoretical models such as the unified fission model (UFM), generalized liquid-drop model (GLDM) and effective liquid-drop model (ELDM). Furthermore, within the CPPM, we predicted the half-lives of potential 2p radioactive nuclei for which experimental data are currently unavailable. The predicted results were then assessed, compared with UFM, ELDM and GLDM models, and examined in detail.

Keywords 2p radioactivity · CPPM · Half-lives · Excited state

1 Introduction

Except for the well-known α , β , γ and cluster decays, etc. [1–10], there exist some exotic modes of radioactivity in proton-rich nuclei [11–13], such as proton and two-proton (2p) radioactivity [14–18]. 2p radioactivity is an extremely exotic mode that is energetically possible in less massive

nuclei near the proton-drip line [19–25], and it can reveal abundant information of nuclear structure, including the nuclear radius, wave function of the emitted two protons, spin and parity, deformation effect and so on [26–28]. This novel decay mode was firstly predicted by Zel'dovich and Goldansky in 1960s [29–31]. Based on the pioneering work of Zel'dovich and Goldansky, an extensive range of theoretical models have been proposed to describe this exotic decay process [32–35]. However, due to the limitations in radioactive beam facilities and detection technology, 2p radioactivity was not experimentally confirmed until the observation of $^{45}\text{Fe} \rightarrow ^{43}\text{Cr} + \text{p} + \text{p}$ decay by Giovinazzo et al. at GANIL (France) and independently by Pfützner et al. at GSI (Germany) in 2002, which provided the first experimental evidence of this decay mode [36, 37]. Since then 2p

This work was supported by the National Natural Science Foundation of China (Nos. 12175100 and 11975132), the Construct Program of the Key Discipline in Hunan Province, the Research Foundation of Education Bureau of Hunan Province, China (No. 18A237), the Natural Science Foundation of Hunan Province, China (No. 2018JJ2321), the Innovation Group of Nuclear and Particle Physics in USC, the Opening Project of Cooperative Innovation Center for Nuclear Fuel Cycle Technology and Equipment, University of South China (No. 2019KFZ10).

✉ Xi-Jun Wu
wuxijun1980@yahoo.cn

✉ Xiao-Hua Li
lixiaohuaphysics@126.com

¹ School of Nuclear Science and Technology, University of South China, Hengyang 421001, China

² Department of Chemical Engineering, Anhui University of Science and Technology, Huainan 232001, China

³ School of Math and Physics, University of South China, Hengyang 421001, China

⁴ College of Physics and Electronics, Central South University, Changsha 410083, China

⁵ National Exemplary Base for International Sci & Tech. Collaboration of Nuclear Energy and Nuclear Safety, University of South China, Hengyang 421001, China

⁶ Cooperative Innovation Center for Nuclear Fuel Cycle Technology and Equipment, University of South China, Hengyang 421001, China

⁷ Key Laboratory of Low Dimensional Quantum Structures and Quantum Control, Hunan Normal University, Changsha 410081, China

radioactivity has been recognized as the significant decay mode for proton-rich nuclei [38], and it has been detected and studied in several nuclei, such as the resonant ground state of ${}^6\text{Be}$ [39, 40] and ${}^{12}\text{O}$ [41], and in the excited state selectively populated in ${}^6\text{Be}$ [40] and ${}^{14}\text{O}$ [42], etc.

Numerous studies have shown that the 2p radioactivity not only occurs from ground state but also from short-lived excited state. Jänecke was the first to discuss the possibility of β -delayed 2p ($\beta 2p$) radioactivity [38], while Goldansky predicted the occurrence of $\beta 2p$ radioactivity could be found in $\beta 2p$ emitters of $Z = 10\text{--}20$. In 1983, Cable et al. reported the first experimental observation of $\beta 2p$ radioactivity [43]. Subsequently, an increasing number of $\beta 2p$ emitters have been detected using silicon detector telescopes, making it possible to measure the energy of two individual protons with high precision. Since Cable et al. [43] discovered the $\beta 2p$ radioactivity from ${}^{22}\text{Al}$, shortly they observed more $\beta 2p$ radioactivity from ${}^{26}\text{P}$ [44] and ${}^{35}\text{Ca}$ [45]. Hereafter, several other $\beta 2p$ nuclei were found, including ${}^{23}\text{Si}$ [46], ${}^{27}\text{S}$ [47], ${}^{31}\text{Ar}$ [48], ${}^{39}\text{Ti}$ [49], ${}^{43}\text{Cr}$ [50, 51] and ${}^{50}\text{Ni}$ [52]. In addition to populating excited state 2p radioactivity via β decays, 2p radioactivity has also been observed from excited state fed by nuclear reactions such as pick up, transfer or fragmentation, including ${}^{14}\text{O}$ [42], ${}^{17,18}\text{Ne}$ [53–57], ${}^{22}\text{Mg}$ [58, 59] and ${}^{28,29}\text{S}$ [60, 61]. In 2006, Mukha et al. first reported 2p radioactivity from ${}^{94}\text{Ag}$ in an experiment at GSI [62].

From a theoretical perspective, several methods have been proposed over the past few decades to study the mechanism of 2p radioactivity, including both microscopic and phenomenological models. In general, there are three distinct ways for proton-rich nuclei to emit two protons: (1) two-body sequential emission, (2) three-body simultaneous emission and (3) diproton emission (also called ${}^2\text{He}$ cluster emission). The ${}^2\text{He}$ cluster emission is an extreme scenario with the two strongly correlated protons that can only survive for a brief time before splitting after passing through the Coulomb barrier. Recently, based on the CPPM, Yao et al. [64] and Ghodsi et al. [65], as well as Deng et al. [66] and Santhosh et al. [67], performed comparative studies of various proximity potential formalisms to study α decay, proton radioactivity and cluster radioactivity, respectively. Considering that the 2p radioactivity decay process shares the same theory as α decay and proton radioactivity, i.e., barrier penetration [68–73], we extended the CPPM presented in Ref. [74] to account for ground state 2p radioactivity. It was found that both the calculated and predicted results were highly consistent with experimental data and results from other theoretical models. In fact, except for theoretical models the empirical formulae are excellent tools for investigating 2p radioactivity involving a four-parameter empirical formula provided by Sreeja et al. [34] and a two-parameter empirical formula proposed by Liu et al. [63]. Consequently, a desirable question is whether 2p radioactivity from excited states can

be regarded as ${}^2\text{He}$ cluster emission and described by the CPPM. To address this question, we systematically studied the half-lives of 2p radioactivity from excited states within the CPPM for nuclei with $8 < Z < 47$ in this work.

This paper is organized as follows: In Sect. 2, the theoretical framework of 2p radioactivity half-life and the CPPM formalism are described briefly. In Sect. 3, detailed calculations and discussion are presented. Finally, a summary is given in Sect. 4.

2 Theoretical framework

2.1 The 2p radioactivity half-life formalism

The half-life of 2p radioactivity is generally determined by

$$T_{1/2} = \frac{\ln 2}{\lambda} = \frac{\ln 2}{\nu P S_{2p}}, \quad (1)$$

where λ is the decay constant. ν is the assault frequency related to the harmonic oscillation frequency presented in the Nilsson potential [75], it can be written as

$$h\nu = \hbar\omega \simeq \frac{41}{A^{1/3}}. \quad (2)$$

Here h , \hbar , A and ω are the Planck constant, reduced Planck constant, mass number of parent nucleus and angular frequency, respectively. $S_{2p} = G^2 [A/(A-2)]^{2n} \chi^2$ represents the preformation probability of the emitted two protons in parent nucleus obtained by the cluster overlap approximation [74]. Here, $G^2 = \frac{(2n)!}{2^{2n}(n!)^2}$ [20, 86] with $n \approx (3Z)^{1/3} - 1$ is the average principal proton oscillator quantum number and χ^2 is set as 0.0143 according to Ref. [23]. The penetration probability P can be calculated by the WKB approximation and written as

$$P = \exp \left[-2 \int_{r_{\text{in}}}^{r_{\text{out}}} K(r) dr \right], \quad (3)$$

where $K(r) = \sqrt{\frac{2\mu}{\hbar^2} |V(r) - Q_{2p}|}$, μ denotes the reduced mass [35]. r is the mass center distance between the daughter nucleus and emitted two protons. r_{in} and r_{out} are classical inner and outer turning points which can be obtained from the conditions $V(r_{\text{in}}) = V(r_{\text{out}}) = Q_{2p}$. The whole interaction potential $V(r)$ between the emitted two protons and daughter nucleus is composed of the Coulomb potential $V_C(r)$, nuclear potential $V_N(r)$ and centrifugal potential $V_\ell(r)$. It can be expressed as

$$V(r) = V_N(r) + V_C(r) + V_\ell(r). \quad (4)$$

In CPPM, the nuclear potential is replaced by proximity potential. A detailed description of the proximity potential will be provided in Sect. 2.2.

Assuming a homogeneous spherical charge distribution for the daughter nucleus, the Coulomb potential $V_C(r)$ is postulated to be the potential of a uniformly charged sphere with radius R . It is expressed as follows:

$$V_C(r) = \begin{cases} \frac{Z_1 Z_2 e^2}{2R} \left[3 - \left(\frac{r}{R}\right) \right], & r < R, \\ \frac{Z_1 Z_2 e^2}{r}, & r > R, \end{cases} \quad (5)$$

where $R = R_1 + R_2$ is the separation radius with R_1 and R_2 being the radii of daughter nucleus and emitted two protons, respectively.

For $V_\ell(r)$, we choose the Langer modified form, because $\ell(\ell + 1) \rightarrow (\ell + \frac{1}{2})^2$ is a necessary correction for one-dimensional problems [76]. It can be written as

$$V_\ell(r) = \frac{\hbar^2(\ell + \frac{1}{2})^2}{2\mu r^2}, \quad (6)$$

where ℓ is the orbital angular momentum taken away by the emitted two protons.

2.2 The proximity potential formalism

The phenomenological proximity potential was first proposed by Blocki et al. [77] in 1970s for heavy-ion reactions. It provides a simple formula for the nucleus–nucleus interaction energy as a function of the separation between the surfaces of the approaching nuclei, with adjustable parameters that makes use of the measured values of the nuclear surface tension and surface diffuseness. In the CPPM, the potential energy barrier is modeled as the sum of Coulomb potential, proximity potential and centrifugal potential for both the touching configuration and separated fragments. In this work, we select the AW95 proposed by Aage Withner in 1995 [78] as an example to replace nuclear potential. This set of proximity potential can be expressed as

$$V_N(r) = -\frac{V_0}{1 + \exp\left(\frac{r-R_0}{0.63}\right)}. \quad (7)$$

Here $V_0 = 16\pi \frac{R_1 R_2}{R_1 + R_2} \gamma a$ with

$$a = \frac{1}{1.17 \left(1 + 0.53 \left(A_1^{-1/3} + A_2^{-1/3} \right) \right)} \text{ fm}. \quad (8)$$

$R_i = (1.2A_i^{1/3} - 0.09)$ fm ($i = 1, 2$) and $R_0 = R_1 + R_2$. The surface energy constant γ has the form as

$$\gamma = \gamma_0 \left[1 - k_s \left(\frac{N_1 - Z_1}{A_1} \right) \left(\frac{N_2 - Z_2}{A_2} \right) \right], \quad (9)$$

where the coefficients $\gamma_0 = 0.95$ MeV/fm² and $k_s = 1.8$. Here A_i , Z_i and N_i ($i = 1, 2$) are the mass number, proton number and neutron number daughter nucleus and emitted two protons, respectively.

3 Results and discussion

Blocki first presented the proximity potential in 1977 to characterize the interaction potential between any two nuclei in the separation degree of freedom, based on the proximity force theorem [77]. Hence, numerous nuclear proximity potentials have been widely used in nuclear physics research [79–83]. The CPPM is a phenomenological model that is commonly used to study the two-body problem in the context of 2p radioactivity. It considers the interaction potential between the parent and daughter nuclei as the sum of the nuclear potential, Coulomb potential and centrifugal potential. Unlike other models, the CPPM utilizes the proximity potential, which replaces the nuclear potential with a simplified formalism based on the proximity force. This provides the model with the advantage of adjustable parameters, making it simple yet accurate. In our previous work [74], we extended the CPPM to study 2p radioactivity from the ground state, the main intention of this work is to further extend the CPPM to investigate 2p radioactivity from excited states.

Firstly, we performed the calculations on 2p radioactivity half-lives for $^{14}\text{O}^*$, $^{17}\text{Ne}^*$, $^{18}\text{Ne}^*$, $^{22}\text{Mg}^*$, $^{29}\text{S}^*$ and $^{94}\text{Ag}^*$ (* represent the excited state), all of the calculated results are listed in Table 1. In this table, the former four columns represent the 2p decay process, experimental 2p radioactivity released energy, spin and parity of the initial and final state of the nucleus, angular momentum taken away by the emitted two protons, respectively. The fifth column is the experimental half-lives of 2p radioactivity of excited state. From the sixth to ninth columns, they represent the logarithmic form of 2p half-lives obtained by CPPM, ELDM, GLDM and UFM, respectively. In general, from this table, it is obviously seen that the theoretical half-lives obtained by CPPM are highly consistent with other theoretical models, except for the nuclei $^{22}\text{Mg}^*$ (with the $Q_{2p} = 6.11$ MeV) and $^{29}\text{S}^*$ (with the $Q_{2p} = 5.12$ MeV). In order to intuitively explain this phenomenon, we plot the interaction potential curve of $^{22}\text{Mg}^*$ and $^{29}\text{S}^*$ in Fig. 1. In Sect. 2.1, the half-lives of 2p radioactivity are depended on penetration probability P which is obtained by Eq. 3 and conditions $V(r_{\text{in}}) = V(r_{\text{out}}) = Q_{2p}$.

Unfortunately, in Fig. 1, the whole interaction potential curve of $^{22}\text{Mg}^*$ and $^{29}\text{S}^*$ can't satisfied the conditions




Fig. 1 (Color online) The whole interaction potential of $^{22}\text{Mg}^*$ and $^{29}\text{S}^*$




Fig. 2 (Color online) Comparing the half-lives of the nuclei which obtained by CPPM in Table 1 with other theoretical models

mentioned above, so that we can't calculate corresponding P , naturally, the half-lives can't be obtained.

For the reason of this phenomenon, we turn our attention to remarkable deformation effect. In fact, the half-lives of 2p radioactivity are highly sensitive to proton–proton interaction due to the pairing effect of valence protons. The quasi-classical ^2He model can't account for the experimentally observed proton–proton correlations, which indicate back-to-back proton emission. Moreover, for the nucleus $^{22}\text{Mg}^*$ and $^{29}\text{S}^*$, their spin-parity of the initial state of the parent nuclei are uncertain, and the values of Q_{2p} are floating, all of these factors have nonnegligible impact on $V(r)$ curve.

In order to describe the agreement of the half-lives for the 2p radioactivity of excited state which calculated by CPPM, partial theoretical results are plotted in Fig. 2. In this figure, the logarithmic half-lives of the 2p radioactivity of excited state

obtained by ELDM, GLDM and UFM are compared with our calculations [84, 85]. For the $^{14}\text{O}^*$, in Table 1, its experimental data are given as $\log_{10}T_{1/2} > -16.12$ s, the calculated half-lives by CPPM are very close to those from ELDM, GLDM and UFM, simultaneously the calculated results are better than from the some empirical formulas such as R-matrix theory [42], Sreeja formula and Liu formula [85], whose half-lives are -18.12 s, -19.94 s and -16.85 s, respectively. In order to intuitively discuss the influence of angular momentum ℓ and release energy Q_{2p} on the half-lives of the 2p radioactivity from excited state, we select $^{14}\text{O}^*$ and $^{94}\text{Ag}^*$ as an example to analyze their contribution to the half-lives. In Table 1, it is obviously found that the half-lives of $^{14}\text{O}^*$ obtained by CPPM for the same value of ℓ nearly have three magnitudes deviation with different released energy Q_{2p} value ($Q_{2p} = 1.20$ MeV and $Q_{2p} = 3.15$ MeV). Simultaneously, for the $^{94}\text{Ag}^*$, identical Q_{2p} but different ℓ also have three magnitudes, while the value of angular momentum are 6–10. In fact, this conclusion is same with our previous work [87]. It is obviously found that either Q_{2p} or ℓ has a nonnegligible contribution to the $^{14}\text{O}^*$ and $^{94}\text{Ag}^*$ for their corresponding theoretical half-lives within CPPM, ELDM and GLDM. The reason for this phenomenon is the theory of these models are similar, the penetration probability P are calculated by WKB approximation. In other word, these theoretical models have a strong dependency with Q_{2p} and ℓ , and we suspect the corresponding experimental data are not precise enough. Configuration mixing and three-body dynamics are commonly utilized in nuclear and molecular physics to describe systems with numerous interacting particles. However, due to the limitations of the phenomenological theoretical model, which fails to account for configuration mixing and three-body dynamics, our calculations treat the two emitted protons as strongly correlated, thus reducing them to a two-body problem. As a result, CPPM cannot provide comprehensive information on particle dynamics. Up to now, the experimental 2p radioactivity data of excited state are rare, the reason is that it is hard to observe this decay process due to the extremely short half-lives. From the scheme of these excited nuclei, it is evident that one-proton (1p) radioactivity also exists alongside two-proton radioactivity during their decay process, as well as three-proton (3p) radioactivity decay process. Therefore, the branching ratios between 1p and 2p decay serve as a crucial index for comprehending the abundant information on nuclear structure of these nuclei and merit further investigation.

4 Conclusion

In this work, we extend the CPPM to study the excited 2p radioactivity of $^{14}\text{O}^*$, $^{17}\text{Ne}^*$, $^{18}\text{Ne}^*$, $^{22}\text{Mg}^*$, $^{29}\text{S}^*$ and $^{94}\text{Ag}^*$. It is found that the theoretical values obtained by CPPM are highly consistent with corresponding experimental data and

Table 1 Comparison between the experimental 2p radioactivity half-lives of excited state and those within different theoretical models

2p emission	Q_{2p} (MeV)	$j_i^\pi \rightarrow j_f^\pi$	ℓ	$\log_{10} T_{1/2}^{\text{Expt.}}$	$\log_{10} T_{1/2}^{\text{CPPM}}$	$\log_{10} T_{1/2}^{\text{ELDM}}$ [84]	$\log_{10} T_{1/2}^{\text{GLDM}}$ [84]	$\log_{10} T_{1/2}^{\text{UFM}}$ [85]
$^{14}\text{O}^* \rightarrow ^{12}\text{C}$	1.20	$2^+ \rightarrow 0^+$	2	> -16.12	-16.65	-15.49	-16.10	-16.02
$^{14}\text{O}^* \rightarrow ^{12}\text{C}$	3.15	$2^+ \rightarrow 0^+$	2		-19.25	-18.22	-19.58	-18.87
$^{14}\text{O}^* \rightarrow ^{12}\text{C}$	3.35	$4^+ \rightarrow 0^+$	4		-17.53	-16.25	-16.76	-15.96
$^{17}\text{Ne}^* \rightarrow ^{15}\text{O}$	0.35	$3/2^- \rightarrow 1/2^-$	2	> -10.59	-8.11	-6.98	-6.79	-7.11
$^{17}\text{Ne}^* \rightarrow ^{15}\text{O}$	0.82	$5/2^- \rightarrow 1/2^-$	2		-13.54	-12.41	-12.68	-12.73
$^{17}\text{Ne}^* \rightarrow ^{15}\text{O}$	0.97	$3/2^- \rightarrow 1/2^-$	1		-15.41	-14.20	-14.68	-14.69
$^{18}\text{Ne}^* \rightarrow ^{16}\text{O}$	0.59	$2^+ \rightarrow 0^+$	2		-11.82	-10.59	-10.96	-10.91
$^{18}\text{Ne}^* \rightarrow ^{16}\text{O}$	1.63	$1^- \rightarrow 0^+$	1	$-16.15^{+0.06}_{-0.06}$	-17.56	-16.34	-17.20	-16.79
$^{22}\text{Mg}^* \rightarrow ^{20}\text{Ne}$	6.11	$- \rightarrow 0^+$	0		-	-19.75	-19.58	-18.97
$^{29}\text{S}^* \rightarrow ^{27}\text{Si}$	1.72–2.52	$- \rightarrow 5/2^+$	0		$-16.851 \sim -14.78$	$-15.5 \sim -13.4$	$-14.7 \sim -17.2$	$-16.4 \sim -14.3$
$^{29}\text{S}^* \rightarrow ^{27}\text{Si}$	4.32–5.12	$- \rightarrow 5/2^+$	0		~ -19.04	$-18.4 \sim -17.8$	$-19.2 \sim -18.8$	$-18.9 \sim -18.5$
$^{94}\text{Ag}^* \rightarrow ^{92}\text{Rh}$	1.90	$21^+ \rightarrow 11^+$	6~10	$1.90^{+0.38}_{-0.20}$	6.25~11.89	9.42~14.63	8.22~13.38	9.38~15.21
$^{94}\text{Ag}^* \rightarrow ^{92}\text{Rh}$	1.98				5.45~11.07	8.61~13.80	7.41~12.55	8.56~14.37
$^{94}\text{Ag}^* \rightarrow ^{92}\text{Rh}$	2.05				4.79~10.39	7.95~13.11	6.74~11.86	7.89~13.68
$^{94}\text{Ag}^* \rightarrow ^{92}\text{Rh}$	3.45				-3.82~1.41	-0.8~4.04	-2.03~2.75	-0.92~4.56

theoretical values obtained by ELDM and GLDM. For the nuclei $^{22}\text{Mg}^*$ and $^{29}\text{S}^*$ (with the $Q_{2p} = 5.12$ MeV), the CPPM is not suitable, the reason for this phenomenon perhaps caused by the angular momenta are not available because the spin-parity of the initial state of the parent nuclei has not determined. In addition, the uncertain values of Q_{2p} also provided strong influence on calculated results. Simultaneously, it is found that the half-lives of excited 2p radioactivity have a strong relationship with Q_{2p} and ℓ . Comparing with the theoretical results obtained by ELDM, GLDM and UFM, the half-lives of the excited 2p radioactivity by CPPM are reliable, it maybe as a positive guideline for future experiments.

Author contributions All authors contributed to the study conception and design. Material preparation, data collection and analysis were performed by Yang-Yang Xu, Li-Jia Chu, Xi-Jun Wu, Biao He and Xiao-Hua Li. The first draft of the manuscript was written by De-Xing Zhu, and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

Data availability The data that support the findings of this study are openly available in Science Data Bank at <https://www.doi.org/10.57760/sciencedb.j00186.00160> and <https://cstr.cn/31253.11.sciencedb.j00186.00160>.

Declarations

Conflict of interest The authors declare that they have no competing interests.

References

1. X.D. Sun, P. Guo, X.H. Li, Systematic study of α decay half-lives for even-even nuclei within a two-potential approach. *Phys. Rev. C* **93**, 034316 (2016). <https://doi.org/10.1103/PhysRevC.93.034316>
2. X.D. Sun, P. Guo, X.H. Li, Systematic study of favored α -decay half-lives of closed shell odd-A and doubly-odd nuclei related to ground and isomeric states. *Phys. Rev. C* **94**, 024338 (2016). <https://doi.org/10.1103/PhysRevC.94.024338>
3. C.Z. Shi, Y.G. Ma, α -clustering effect on flows of direct photons in heavy-ion collisions. *Nucl. Sci. Tech.* **32**, 66 (2021). <https://doi.org/10.1007/s41365-021-00897-9>
4. M. Ji, C. Xu, Quantum anti-zeno effect in nuclear decay. *Chin. Phys. Lett.* **38**, 032301 (2021). <https://doi.org/10.1088/0256-307X/38/3/032301>
5. C.W. Ma, H.L. Wei, X.Q. Liu et al., Nuclear fragments in projectile fragmentation reactions. *Prog. Part. Nucl. Phys.* **121**, 103911 (2021). <https://doi.org/10.1016/j.pnpnp.2021.103911>
6. C.W. Ma, J.P. Wei, X.X. Chen et al., Precise machine learning models for fragment production in projectile fragmentation reactions using Bayesian neural networks. *Chin. Phys. C* **46**, 074104 (2022). <https://doi.org/10.1088/1674-1137/ac5efb>
7. L.L. Zhu, B. Wang, M. Wang, H. Zheng, Energy and centrality dependence of light nuclei production in relativistic heavy-ion collisions. *Nucl. Sci. Tech.* **33**, 45 (2022). <https://doi.org/10.1007/s41365-022-01028-8>
8. C. Shen, L. Yan, Recent development of hydrodynamic modeling in heavy-ion collisions. *Nucl. Sci. Tech.* **31**, 122 (2020). <https://doi.org/10.1007/s41365-020-00829-z>
9. F. Zhang, J. Su, Probing neutron-proton effective mass splitting using nuclear stopping and isospin mix in heavy-ion collisions in

- GeV energy region. Nucl. Sci. Tech. **31**, 77 (2020). <https://doi.org/10.1007/s41365-020-00787-6>
10. Y.J. Wang, F.H. Guan, X.Y. Diao et al., CSHINE for studies of HBT correlation in heavy ion reactions. Nucl. Sci. Tech. **32**, 4 (2021). <https://doi.org/10.1007/s41365-020-00842-2>
 11. M. Pfützner, M. Karny, L.V. Grigorenko et al., Radioactive decays at limits of nuclear stability. Rev. Mod. Phys. **84**, 567 (2012). <https://doi.org/10.1103/RevModPhys.84.567>
 12. E. Olsen, M. Pfützner, N. Birge, Landscape of two-proton radioactivity. Phys. Rev. Lett. **110**, 222501 (2013). <https://doi.org/10.1103/PhysRevLett.110.222501>
 13. K.W. Brown, R.J. Charity, L.G. Sobotka et al., Observation of long-range three-body coulomb effects in the decay of ^{16}Ne . Phys. Rev. Lett. **113**, 232501 (2014). <https://doi.org/10.1103/PhysRevLett.113.232501>
 14. K.P. Santhosh, Indu Sukumaran, Description of proton radioactivity using the Coulomb and proximity potential model for deformed nuclei. Phys. Rev. C **96**, 034619 (2017). <https://doi.org/10.1103/PhysRevC.96.034619>
 15. D.S. Delion, S.A. Ghinescu, Two-proton emission systematics. Phys. Rev. C **105**, L031301 (2022). <https://doi.org/10.1103/PhysRevC.105.L031301>
 16. D.N. Basu, P. Roy Chowdhury, C. Samanta, Folding model analysis of proton radioactivity of spherical proton emitters. Phys. Rev. C **72**, 051601(R) (2005). <https://doi.org/10.1103/PhysRevC.72.051601>
 17. X. Pan, Y.T. Zou, H.M. Liu et al., Released energy formula for proton radioactivity based on the liquid-drop model. Commun. Theor. Phys. **73**, 075302 (2021). <https://doi.org/10.1088/1572-9494/abf822>
 18. H.M. Liu, J. Zhang, Z.H. Li et al., Microscopic nuclear equation of state at finite temperature and stellar stability. Phys. Rev. C **106**, 025801 (2022). <https://doi.org/10.1103/PhysRevC.106.025801>
 19. Y.T. Wang, D.Q. Fang, X.X. Xu et al., Implantation-decay method to study the β -delayed charged particle decay. Nucl. Sci. Tech. **29**, 98 (2018). <https://doi.org/10.1007/s41365-018-0438-5>
 20. B.A. Brown, Diproton decay of nuclei on the proton drip line. Phys. Rev. C **43**, R1513 (1991). <https://doi.org/10.1103/PhysRevC.43.R1513>
 21. L.V. Grigorenko, R.C. Johnson, I.G. Mukha et al., Theory of two-proton radioactivity with application to ^{19}Mg and ^{48}Ni . Phys. Rev. Lett. **85**, 22 (2000). <https://doi.org/10.1103/PhysRevLett.85.22>
 22. I. Mukha, L.V. Grigorenko, X. Xu et al., Observation and spectroscopy of new proton-unbound isotopes ^{30}Ar and ^{29}Cl : an interplay of prompt two-proton and sequential decay. Phys. Rev. Lett. **115**, 202501 (2015). <https://doi.org/10.1103/PhysRevLett.115.202501>
 23. J.P. Cui, Y.H. Gao, Y.Z. Wang et al., Two-proton radioactivity within a generalized liquid drop model. Phys. Rev. C **101**, 014301 (2020). <https://doi.org/10.1103/PhysRevC.101.014301>
 24. G. Royer, Calculation of two-proton radioactivity and application to ^9Be , ^6Li , ^3He , and ^2H emissions. Phys. Rev. C **106**, 034605 (2022). <https://doi.org/10.1103/PhysRevC.106.034605>
 25. F.Z. Xing, J.P. Cui, Y.H. Gao, J.Z. Gu, Two-proton radioactivity of exotic nuclei beyond proton drip-line. Commun. Theor. Phys. **73**, 075301 (2021). <https://doi.org/10.1088/1572-9494/abfa00>
 26. B. Blank, M. Płoszajczak, Two-proton radioactivity. Rep. Prog. Phys. **71**, 046301 (2008). <https://doi.org/10.1088/0034-4885/71/4/046301>
 27. A. Kruppa, W. Nazarewicz, Gamow and R-matrix approach to proton emitting nuclei. Phys. Rev. C **69**, 054311 (2004). <https://doi.org/10.1103/PhysRevC.69.054311>
 28. L.S. Ferreira, M.C. Lopea, E. Maglione, Decays of drip line nuclei. Prog. Part. Nucl. Phys. **59**, 418 (2007). <https://doi.org/10.1016/j.pnpnp.2007.01.011>
 29. V.I. Goldansky, On neutron-deficient isotopes of light nuclei and the phenomena of proton and two-proton radioactivity. Nucl. Phys. **19**, 482 (1960). [https://doi.org/10.1016/0029-5582\(60\)90258-3](https://doi.org/10.1016/0029-5582(60)90258-3)
 30. V.I. Goldansky, Two-proton radioactivity. Nucl. Phys. **27**, 648 (1961). [https://doi.org/10.1016/0029-5582\(61\)90309-1](https://doi.org/10.1016/0029-5582(61)90309-1)
 31. Y.B. Zel'dovich, On the existence of new isotopes of light nuclei and the equation of state of neutrons. Sov. Phys. JETP **11**, 812 (1960)
 32. B.A. Brown, B. Blank, J. Giovinazzo, Hybrid model for two-proton radioactivity. Phys. Rev. C **100**, 054332 (2019). <https://doi.org/10.1103/PhysRevC.100.054332>
 33. O.A.P. Tavares, E.L. Medeiros, A calculation model to half-life estimate of two-proton radioactive decay process. Eur. Phys. J. A **54**, 65 (2018). <https://doi.org/10.1140/epja/i2018-12495-4>
 34. I. Sreeja, M. Balasubramaniam, An empirical formula for the half-lives of exotic two-proton emission. Eur. Phys. J. A **55**, 33 (2019). <https://doi.org/10.1140/epja/i2019-12694-5>
 35. H.M. Liu, X. Pan, Y.T. Zou et al., Systematic study of two-proton radioactivity within a Gamow-like model. Chin. Phys. C **45**, 044110 (2021). <https://doi.org/10.1088/1674-1137/abe10f>
 36. M. Pfützner, E. Badura, C. Bingham et al., First evidence for the two-proton decay of ^{45}Fe . Eur. Phys. J. A **14**, 279 (2002). <https://doi.org/10.1140/epja/i2002-10033-9>
 37. J. Giovinazzo, B. Blank, M. Chartier et al., Two-proton radioactivity of ^{45}Fe . Phys. Rev. Lett. **89**, 102501 (2002). <https://doi.org/10.1103/PhysRevLett.89.102501>
 38. J. Jänecke, The emission of protons from light neutron-deficient nuclei. Nucl. Phys. **61**, 326 (1965). [https://doi.org/10.1016/0029-5582\(65\)90907-7](https://doi.org/10.1016/0029-5582(65)90907-7)
 39. D.F. Geesaman, R.L. McGrath, P.M.S. Lesser et al., Particle decay of ^6Be . Phys. Rev. C **15**, 1835 (1977). <https://doi.org/10.1103/PhysRevC.15.1835>
 40. O.V. Bochkarev, L.V. Chulkov, A.A. Korshennikov et al., Democratic decay of ^6Be states. Nucl. Phys. A **505**, 215 (1989). [https://doi.org/10.1016/0375-9474\(89\)90371-0](https://doi.org/10.1016/0375-9474(89)90371-0)
 41. R.A. Kryger, A. Azhari, M. Hellström et al., Two-Proton Emission from the Ground State of ^{12}O . Phys. Rev. Lett. **74**, 860 (1995). <https://doi.org/10.1103/PhysRevLett.74.860>
 42. C.R. Bain, P.J. Woods, R. Coszach et al., Two proton emission induced via a resonance reaction. Phys. Lett. B **373**, 35 (1996). [https://doi.org/10.1016/0370-2693\(96\)00109-8](https://doi.org/10.1016/0370-2693(96)00109-8)
 43. M.D. Cable, J. Honkanen, R.F. Parry et al., Discovery of beta-delayed two-proton radioactivity: ^{22}Al . Phys. Rev. Lett. **50**, 404 (1983). <https://doi.org/10.1103/PhysRevLett.50.404>
 44. J. Honkanen, M.D. Cable, R.F. Parry et al., Beta-delayed two-proton decay of ^{26}P . Phys. Lett. B **133**, 146 (1983). [https://doi.org/10.1016/0370-2693\(83\)90547-6](https://doi.org/10.1016/0370-2693(83)90547-6)
 45. J. Äystö, D.M. Moltz, X.J. Xu et al., Phys. Rev. Lett. **55**, 1384 (1985). <https://doi.org/10.1103/PhysRevLett.55.1384>
 46. B. Blank, F. Boué, S. Andriamonje et al., Spectroscopic studies of the βp and $\beta 2p$ decay of ^{23}Si . Z. Phys. A **357**, 247 (1997). <https://doi.org/10.1007/s002180050241>
 47. V. Borrel, J.C. Jacmart, F. Pougheon, ^{31}Ar and ^{27}S : beta-delayed two-proton emission and mass excess. Nucl. Phys. **531**, 353 (1991). [https://doi.org/10.1016/0375-9474\(91\)90616-E](https://doi.org/10.1016/0375-9474(91)90616-E)
 48. J.E. Reiff, M.A.C. Hotchkiss, D.M. Moltz, A fast in-beam recoil catcher wheel and the observation of beta-delayed two-proton emission from ^{31}Ar . Nucl. Instrum. Methods. A **276**, 228 (1989). [https://doi.org/10.1016/0168-9002\(89\)90637-2](https://doi.org/10.1016/0168-9002(89)90637-2)
 49. D.M. Moltz, J.C. Batchelder, T.F. Lang et al., Beta-delayed two-proton decay of ^{39}Ti . Z. Phys. A **342**, 273 (1992). <https://doi.org/10.1007/BF01291509>
 50. V. Borrel, R. Anne, D. Bazin et al., The decay modes of proton drip-line nuclei with A between 42 and 47. Z. Phys. A **344**, 135 (1992). <https://doi.org/10.1007/BF01291696>

51. J. Giovinazzo, B. Blank, C. Borcea et al., First direct observation of two protons in the decay of ^{45}Fe with a time-projection chamber. *Phys. Rev. Lett.* **99**, 102501 (2007). <https://doi.org/10.1103/PhysRevLett.99.102501>
52. C. Dossat, N. Adimi, F. Aksouh et al., The decay of proton-rich nuclei in the mass $A = 36 - 56$ region. *Nucl. Phys. A* **792**, 18 (2007). <https://doi.org/10.1016/j.nuclphysa.2007.05.004>
53. M.J. Chromik, B.A. Brown, M. Fauerbach et al., Excitation and decay of the first excited state of ^{17}Ne . *Phys. Rev. C* **55**, 1676 (1997). <https://doi.org/10.1103/PhysRevC.55.1676>
54. J. Gómez del Campo, A. Galindo-Uribarri, J.R. Beene et al., Decay of a resonance in by the simultaneous emission of two protons. *Phys. Rev. Lett.* **86**, 43 (2001). <https://doi.org/10.1103/PhysRevLett.86.43>
55. G. Raciti, G. Cardella, M. De Napoli et al., Experimental evidence of ^2He decay from ^{18}Ne excited states. *Phys. Rev. Lett.* **100**, 192503 (2008). <https://doi.org/10.1103/PhysRevLett.100.192503>
56. M.J. Chromik, P.G. Thierolf, M. Thoennessen et al., Two-proton spectroscopy of low-lying states in ^{17}Ne . *Phys. Rev. C* **66**, 024313 (2002). <https://doi.org/10.1103/PhysRevC.66.024313>
57. T. Zerguerras, B. Blank, Y. Blumenfeld et al., Study of light proton-rich nuclei by complete kinematics measurements. *Eur. Phys. J. A* **20**, 389 (2004). <https://doi.org/10.1140/epja/i2003-10176-1>
58. Y.G. Ma, D.Q. Fang, X.Y. Sun et al., Different mechanism of two-proton emission from proton-rich nuclei ^{23}Al and ^{22}Mg . *Phys. Lett. B* **743**, 306 (2015). <https://doi.org/10.1016/j.physletb.2015.02.066>
59. D.Q. Fang, Y.G. Ma, X.Y. Sun et al., Proton-proton correlations in distinguishing the two-proton emission mechanism of ^{23}Al and ^{22}Mg . *Phys. Rev. C* **94**, 044621 (2016). <https://doi.org/10.1103/PhysRevC.94.044621>
60. C.J. Lin, X.X. Xu, H.M. Jia et al., Experimental study of two-proton correlated emission from ^{29}S excited states. *Phys. Rev. C* **80**, 014310 (2009). <https://doi.org/10.1103/PhysRevC.80.014310>
61. X.X. Xu, C.J. Lin, H.M. Jia et al., Correlations of two protons emitted from excited states of ^{28}S and ^{27}P . *Phys. Lett. B* **727**, 126 (2013). <https://doi.org/10.1016/j.physletb.2013.10.029>
62. I. Mukha, E. Roeckl, L. Batist et al., Proton-proton correlations observed in two-proton radioactivity of ^{94}Ag . *Nature* **439**, 298 (2006). <https://doi.org/10.1038/nature04453>
63. H.M. Liu, Y.T. Zou, X. Pan et al., New Geiger-Nuttall law for two-proton radioactivity. *Chin. Phys. C* **45**, 024108 (2021). <https://doi.org/10.1088/1674-1137/abd01e>
64. Y.J. Yao, G.L. Zhang, W.W. Qu et al., Comparative studies for different proximity potentials applied to α decay. *Eur. Phys. J. A* **51**, 122 (2015). <https://doi.org/10.1140/epja/i2015-15122-0>
65. O.N. Ghodsi, A. Daei-Ataollah, Systematic study of α decay using various versions of the proximity formalism. *Phys. Rev. C* **93**, 024612 (2016). <https://doi.org/10.1103/PhysRevC.93.024612>
66. J.G. Deng, X.H. Li, J.L. Chen et al., Systematic study of proton radioactivity of spherical proton emitters within various versions of proximity potential formalisms. *Eur. Phys. J. A* **55**, 58 (2019). <https://doi.org/10.1140/epja/i2019-12728-0>
67. K.P. Santhosh, I. Sukumaran, Heavy particle decay studies using different versions of nuclear potentials. *Eur. Phys. J. Plus* **132**, 431 (2019). <https://doi.org/10.1140/epjp/i2017-11743-x>
68. Y.Z. Wang, S.J. Wang, Z.Y. Hou, J.Z. Gu, Systematic study of α -decay energies and half-lives of superheavy nuclei. *Phys. Rev. C* **92**, 064301 (2015). <https://doi.org/10.1103/PhysRevC.92.064301>
69. Y.Z. Wang, J.P. Cui, Y.L. Zhang et al., Competition between α decay and proton radioactivity of neutron-deficient nuclei. *Phys. Rev. C* **92**, 064301 (2017). <https://doi.org/10.1103/PhysRevC.95.014302>
70. Y.Z. Wang, F.Z. Xing, Y. Xiao, J.Z. Gu, An improved semi-empirical relationship for cluster radioactivity. *Chin. Phys. C* **45**, 044111 (2021). <https://doi.org/10.1088/1674-1137/abe112>
71. J.G. Deng, J.H. Cheng, B. Zheng, X.H. Li, α decay properties of ^{297}Og within the two-potential approach. *Chin. Phys. C* **41**, 124109 (2017). <https://doi.org/10.1088/1674-1137/41/12/124109>
72. H.M. Liu, Y.T. Zou, P. Xiao et al., Systematic study of the α decay preformation factors of the nuclei around the $Z = 82, N = 126$ shell closures within the generalized liquid drop model. *Chin. Phys. C* **44**, 094106 (2020). <https://doi.org/10.1088/1674-1137/44/9/094106>
73. F.Z. Xing, H. Qi, J.P. Cui et al., An improved Gamow-like formula for α -decay half-lives. *Nucl. Phys. A* **1028**, 122528 (2022). <https://doi.org/10.1016/j.nuclphysa.2022.122528>
74. D.X. Zhu, H.M. Liu, Y.Y. Xu et al., Two-proton radioactivity within Coulomb and proximity potential model. *Chin. Phys. C* **46**, 044106 (2022). <https://doi.org/10.1088/1674-1137/ac45ef>
75. S.G. Nilsson, Binding states of individual nucleons in strongly deformed nuclei. *Dan. Mat. Fys. Medd.* **29**, 16 (1955)
76. J.J. Morehead, Asymptotics of radial wave equations. *J. Math. Phys.* **36**, 5431 (1955). <https://doi.org/10.1063/1.531270>
77. J. Blocki, J. Randrup, W.J. Świątecki et al., Proximity forces. *Ann. Phys.* **105**, 427 (1977). [https://doi.org/10.1016/0003-4916\(77\)90249-4](https://doi.org/10.1016/0003-4916(77)90249-4)
78. A. Winther, Dissipation, polarization and fluctuation in grazing heavy-ion collisions and the boundary to the chaotic regime. *Nucl. Phys. A* **594**, 203 (1995). [https://doi.org/10.1016/0375-9474\(95\)00374-A](https://doi.org/10.1016/0375-9474(95)00374-A)
79. C.L. Guo, G.L. Zhang, Analysis of proton radioactivity of nuclei by using proximity potential with a new universal function. *Eur. Phys. J. A* **50**, 187 (2014). <https://doi.org/10.1140/epja/i2014-14187-5>
80. L. Zheng, G.L. Zhang et al., Calculation of half-lives of cluster decay by using proximity potential with a new universal function. *Nucl. Phys. A* **915**, 70 (2013). <https://doi.org/10.1016/j.nuclphysa.2013.06.012>
81. W.W. Qu, G.L. Zhang et al., Comparative studies of Coulomb barrier heights for nuclear models applied to sub-barrier fusion. *Phys. Rev. C* **90**, 064603 (2014). <https://doi.org/10.1103/PhysRevC.90.064603>
82. R. Kumar, M.K. Sharma et al., Systematic study of various proximity potentials in ^{208}Pb -daughter cluster radioactivity. *Phys. Rev. C* **85**, 054612 (2014). <https://doi.org/10.1103/PhysRevC.85.054612>
83. I. Dutt, R.K. Puri, Comparison of different proximity potentials for asymmetric colliding nuclei. *Phys. Rev. C* **81**, 064609 (2010). <https://doi.org/10.1103/PhysRevC.81.064609>
84. F.Z. Xing, J.P. Cui, Y.Z. Wang et al., Two-proton emission from excited states of proton-rich nuclei. *Acta Phys. Sin.* **71**, 062301 (2022). <https://doi.org/10.7498/aps.71.20211839>
85. F.Z. Xing, J.P. Cui, Y.Z. Wang, J.Z. Gu, Two-proton radioactivity of ground and excited states within a unified fission model. *Chin. Phys. C* **45**, 124105 (2021). <https://doi.org/10.1088/1674-1137/ac2425>
86. N. Anyas-Weiss, J.C. Cornell, P.S. Fisher et al., Nuclear structure of light nuclei using the selectivity of high energy transfer reactions with heavy ions. *Phys. Rep.* **12**, 101 (1974). [https://doi.org/10.1016/0370-1573\(74\)90045-3](https://doi.org/10.1016/0370-1573(74)90045-3)
87. D.X. Zhu, Y.Y. Xu, H.M. Liu et al., ITwo-proton radioactivity of the excited state within the Gamowlike and modified Gamow-like models. *Nucl. Sci. Tech.* **32**, 122 (2022). <https://doi.org/10.1007/s41365-022-01116-9>

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.