

# Gamma-, neutron-, and muon-induced environmental background simulations for <sup>100</sup>Mo-based bolometric double-beta decay experiment at Jinping Underground Laboratory

Wei Chen<sup>1</sup> · Long Ma<sup>1</sup> · Jin-Hui Chen<sup>1</sup> · Huan-Zhong Huang<sup>1,2</sup>

Received: 8 May 2023 / Revised: 21 July 2023 / Accepted: 30 July 2023 / Published online: 20 September 2023 © The Author(s), under exclusive licence to China Science Publishing & Media Ltd. (Science Press), Shanghai Institute of Applied Physics, the Chinese Academy of Sciences, Chinese Nuclear Society 2023

## Abstract

The sensitivity of an experiment to detect the Majorana neutrino mass via neutrinoless double-beta decay  $(0\nu\beta\beta)$  strongly depends on the rate of background events that can mimic this decay. One major source of this background is the radioactive emissions from the laboratory environment. In our study, we focused on assessing the background contributions from environmental gamma rays, neutrons, and underground muons to the Jinping bolometric demonstration experiment. This experiment uses an array of lithium molybdate crystal bolometers to probe the potential  $0\nu\beta\beta$  decay of the <sup>100</sup>Mo isotope at the China Jinping Underground Laboratory. We also evaluated the shielding effectiveness of the experimental setup through an attenuation study. Our simulations indicate that the combined background from environmental gamma rays, neutrons, and muons in the relevant <sup>100</sup>Mo  $0\nu\beta\beta$  *Q*-value region can be reduced to approximately 0.003 cts/kg/keV/yr.

Keywords Neutrinoless double-beta decay · Geant4 · CUPID

# 1 Introduction

Unveiling the properties of neutrinos is fundamental, and searching for neutrinoless double-beta  $(0\nu\beta\beta)$  decay stands as a pivotal approach to this goal. Observing the  $0\nu\beta\beta$  decay experimentally can directly substantiate the Majorana nature of neutrinos, implying that neutrinos are their own antiparticles. This would open new frontiers in physics, extending beyond the standard framework of particle physics [1].

This work was supported in part by the State Key Research Development Program in China (Nos. 2022YFA1604702 and 2022YFA1604900), the National Natural Science Foundation of China (No. 12025501), and Strategic Priority Research Program of Chinese Academy of Science (No. XDB34030200).

 Long Ma malong@fudan.edu.cn
 Jin-Hui Chen chenjinhui@fudan.edu.cn

<sup>1</sup> Key Laboratory of Nuclear Physics and Ion-Beam Application (MOE), Institute of Modern Physics, Fudan University, Shanghai 200433, China

<sup>2</sup> University of California, Los Angeles, CA 90095, USA

Numerous experimental techniques have been employed to detect the rare  $0\nu\beta\beta$  events. While direct evidence of  $0\nu\beta\beta$ remains elusive, strict limits on the half-life of various double-beta decay nuclides have been established [2–5]. Among the tools used, crystal bolometers, which operate at ultralow temperatures, stand out. They are preferred for  $0\nu\beta\beta$ searches due to their high energy resolution, efficient detection, and stable operation [6]. These cryogenic bolometers employ crystals embedded with double-beta decay isotopes to detect the  $0\nu\beta\beta$  decay. The forthcoming CUPID experiment (CUORE with particle identification), integrating a high-performance heat-light dual readout to discern alpha particle backgrounds, is projected to enhance the sensitivity of  $0\nu\beta\beta$  detection in future [7–9].

A critical challenge in  $0\nu\beta\beta$  detection is the potential interference from external radioactivity, as it can significantly hamper the experiment's potential. Efforts have thus been made to understand and mitigate sources of background interference. Previous research has identified multiple culprits for misleading counts in the  $0\nu\beta\beta$  search energy region of interest (ROI), including cosmogenic activation, natural radioactive contamination from materials, and environmental radioactive emissions in the laboratory [10–14]. Given that cosmogenic activation at sea level notably influences the detectors, experiments are usually conducted in deep mines or under mountains; these locations, shielded by the earth's crust or mountain rock, offer protection from cosmic radiation [15, 16]. Besides residual cosmogenic background, environmental radioactive interference remains crucial for underground setups. Main sources include gamma rays from rocks, neutrons from surrounding materials, and events induced by underground muons [17, 18]. Passive shields are typically deployed to counter gamma and neutron backgrounds, while muon vetoes are used to effectively tag and exclude underground muon events and their induced activities.

Given that background levels are pivotal for experimental sensitivity, thorough studies on environmental background shielding and subsequent evaluations of any remaining background are crucial during the design phase of the detector system. In this paper, leveraging a Geant4-based Monte Carlo simulation, we delve into the background contributions-due to environmental gamma rays, neutrons, and cosmic muons-to the Jinping CUPID demonstration experiment. This experiment, featuring a 10-kg cryogenic crystal bolometer at the China Jinping Underground Laboratory (CJPL), aims to showcase the advanced technologies intended for the next-generation bolometric experiment targeting the  $0\nu\beta\beta$  decay of <sup>100</sup>Mo. This paper is structured as follows: The simulation setup and methods are discussed in Sect. 2. Findings related to particle attenuation and background are elaborated in Sect. 3. Conclusions and discussion are given in Sect. 4

## 2 Simulation setup

We utilized the Geant4 simulation toolkit (version 4.10.5) to model radiation shielding and assess background contributions [19]. Geant4 is renowned for offering an extensive array of particle-material interaction cross sections, allowing precise characterization of particle production and transport properties [20-22]. For our study, we employed the "Shielding" physics list, which is tailored for simulations associated with deep underground rare-event experiments. This physics list incorporates a series of interaction models, each detailing specific physical processes. For neutrons with energies up to 20 MeV, we leveraged high-precision models and cross sections to depict processes such as elastic and inelastic scattering, capture, and fission. To study muoninduced backgrounds, we also applied the "QGSP\_BERT" physics list as a supplementary check, given its suitability for cosmic ray-related processes. Event generation for gamma rays and neutrons was based on the in situ spectral measurements taken in the CJPL experimental hall, as depicted in Fig. 1. We assumed isotropic radiation for both gamma and neutron sources. The initiation points for these environmental background sources were set in line with the hall's dimensions (14 m in height and 12 m in width). Moreover, the muon simulation factored in experimentally measured zenith and azimuthal angle distributions.

Figure 2 shows a schematic view of the experimental setup, including the detector crystal array, background shields surrounding the detector, and cooling system used to maintain low temperatures. The configurations of the detector and shielding systems are implemented in the simulation code. The CUPID-CJPL demonstrator comprised an array of 36 lithium molyb-date ( $Li_2MoO_4$ , abbr. LMO) cryogenic bolometers, having a cumulative crystal mass of approximately 10 kg. The LMO



Fig. 1 (Color online) Energy spectra of CJPL environmental gamma (left) and neutron (right). The sampled Monte Carlo spectra (solid lines) are shown in comparison with the results obtained by the in situ measurements (dashed lines) [23, 28]





Table 1 Key elements of the experimental setup

Element	Material	Mass (kg)	Surface (cm <sup>2</sup> )
Crystal array	LMO	10.0	4374.0
Thermal shield	Cu-OFE	61.7	23850.9
Top Cu shield	Cu-OFE	209.7	4736.9
Lateral Cu shield	Cu-OF	2990.4	60880.0
Lateral PE shield	PE	1343.9	114300.0

crystals not only serve as the source for the  $0\nu\beta\beta$  decay but also function as heat absorbers. This array is organized into nine layers, with each layer containing four bolometers. Each of these bolometers is cooled to an ultra-low temperature of about 10 mK using a dilution refrigerator. The individual bolometer module integrates a 4.5 cm × 4.5 cm × 4.5 cm LMO cubic crystal and is equipped with a heat-light dual-phase readout. For these modules, 280 g LMO crystals were specifically chosen. The cooling apparatus, or the cryostat, consists of three concentric vessels dedicated to thermal shielding. To complement these thermal barriers, radiation shields were strategically positioned both inside and surrounding the cryostat to counteract radiation emanating from the cryogenic system. Within this arrangement, a substantial copper shield (having a thickness of 12 cm and a diameter of 50 cm) was situated above the detector tower to mitigate external radioactive interference. To further fortify the system, passive shielding materials were employed. This includes a 12-cm-thick layer of copper, designed to diminish environmental gamma rays, and a 20-cm-thick layer of 5%-borated polyethylene (PE) tailored to intercept environmental neutrons. The salient features of this experimental setup are detailed in Table 1.

# **3** Results and discussion

## 3.1 Attenuation study

Numerous experimental studies have established that materials with high atomic mass numbers, such as lead and copper, excel at attenuating gamma rays. Notably, copper typically possesses lower inherent radioactivity than lead. As a result, high-purity oxygen-free copper has emerged as a preferred material for crafting shielding systems in sensitive underground experiments related to rare events.

For the Jinping bolometric demonstration, the design of the background shielding system was informed by a Monte Carlo simulation. Utilizing the Geant4 toolkit, we simulated particle interactions with materials to quantitatively assess the attenuation capacities of the shielding against gamma rays and neutrons.

Figure 3 shows the gamma-ray absorption fraction as a function of the gamma energy for different copper shielding thicknesses. The incident gamma energy was sampled based on the experimentally measured CJPL in situ gamma spectra (Fig 1). Our results indicate that the gamma absorption at different energies improves significantly by increasing the thickness of the material. In addition to high-energy environmental gamma rays, fast neutrons with energies above 1 MeV are serious background sources for rare underground event experiments. Borated polyethylene (BPE) is an effective material for fast neutron shielding [24, 25]. The results of the neutronshielding performance are shown in Fig. 4 compared with the gamma results. The black and blue curves represent the gamma and neutron absorptions of copper and 5% BPE, respectively. It was observed that the neutron shielding



**Fig.3** (Color online) Simulated results of the gamma absorption for varying thicknesses of the copper brick



**Fig. 4** (Color online) Simulated results of the attenuation coefficient as a function of copper shielding thickness for gamma and borated polyethylene thickness for neutron. The input energy spectra for gamma and neutron are shown in Fig. 1

was improved by increasing the thickness of BPE. A reduction in gamma and neutron fluxes by a factor of  $\sim 10$  can be realized by deploying a 12-cm-thick copper shield and an 8-cm 5% BPE shield, respectively. Thus, for the CJPL bolometric experiment, with 12-cm-thick copper and 20-cm-thick 5%-borated polyethylene external shields, we expect a reduction exceeding tenfold in the environmental gamma and neutron fluxes within the detector system.

#### 3.2 Environmental gamma-induced background

Different background sources have the potential to simulate a  $0\nu\beta\beta$  signal within the decay-energy ROI of  $0\nu\beta\beta$ . A predominant source of such radiation is environmental gamma rays. For underground experiments, these gamma rays largely stem from the decay of radioactive nuclei and processes induced by high-energy neutrons. Moreover, the lab's infrastructure and its experimental apparatus can introduce additional gamma background. In our research, we delved into evaluating the contribution of environmental gamma background to a <sup>100</sup>Mo  $0\nu\beta\beta$  detection using a bolometric method.

The gamma radioactivity of the CJPL environment has been extensively profiled in prior studies through in situ gamma-ray measurements employing a high-purity germanium spectrometer [26–30]. The naturally occurring gamma spectrum spans up to 3 MeV; events exceeding this energy are typically the result of interactions between neutrons, muons, and rocks. Given that the isotope <sup>100</sup>Mo boasts a relatively high  $Q_{\beta\beta}$  value (approximately 3034.40(17) keV), we designated the ROI to range from 3000 to 3060 keV. This decision factored in the energy resolution of the crystal bolometer when assessing the background. The approach to parameterizing the energy resolution is as follows:

$$\sigma(E) = \sqrt{0.49 + (0.058 \times \sqrt{E})^2},$$
(1)

where E denotes the measured energy in keV based on the CUPID-Mo study [31].

The gamma spectrum measured using a high-purity germanium (HPGe) detector in CJPL Phase II was used as the input for our Geant4 simulation [26]. In this study, given that we focused on the <sup>100</sup>Mo  $0\nu\beta\beta$  ROI, only high-energy gamma rays ( $E_{\gamma} > 3000$  keV) were considered. The integral flux of the high-energy gamma rays was calculated as  $2.51 \pm$  $0.07 \times 10^{-6}$  cm<sup>-2</sup>s<sup>-1</sup> based on the experimentally measured results.

In bolometric experiments, the array structure of the detector system allows us to apply the anti-coincidence cut criterion. This helps in rejecting high-energy external back-ground events that pass through several crystal modules. Figure 5 illustrates the concept of "*multiplicity*". An event that deposits energy in a single crystal module is defined as having "*multiplicity* = 1". Conversely, if the energy is deposited across multiple crystal modules, it is classified as a "*multiplicity* > 1" event. By selecting the "*multiplicity* = 1" events, we can significantly filter out high-energy external gamma rays, as well as neutrons and muons, that traverse multiple crystals.

Figure 6 presents the simulated results for the gammainduced background. The results for different cutting criteria are shown for comparison. It is observed that by requiring



Fig. 5 (Color online) Definition of the multiplicity



Fig. 6 (Color online) Simulated results of the gamma-induced background

"multiplicity = 1", the environmental gammas contribute approximately  $1.3 \times 10^{-3}$  cts/keV/kg/yr background in the <sup>100</sup>Mo  $0\nu\beta\beta$  ROI (3000–3060 keV). The uncertainty of the result is estimated by considering the statistical uncertainties of the flux measurements and selection of PhysicsList ("Shielding" vs "QGSP\_BERT\_HP"). The relative difference was found to be within 10%.

#### 3.3 Environmental neutron-induced background

High-energy neutrons present another significant background for  $0\nu\beta\beta$  experiments. In deep underground laboratories, neutron fluxes below 10 MeV primarily arise from spontaneous fission (such as from U-238) and ( $\alpha$ ,*n*) reactions, which involve interactions between alphas from natural radioactive sources and light target nuclei within the rock. Beyond production by the inherent radioactivity of materials, neutrons can also be produced by cosmic ray muons and their secondary particles through nuclear reactions [32, 33]. The energy of these muon-induced neutrons can reach up to several gigaelectronvolts.

Systematic measurements of the fluxes of low-energy thermal neutrons and high-energy fast neutrons have been performed in deep underground laboratories [23, 34–36]. In our simulation study, we evaluated the neutron-induced background based on the in situ fast neutron spectrum measured at the CJPL Hall-A [37]. Monte Carlo neutrons with kinetic energies exceeding 1 MeV were sampled based on the experimental flux spectrum results. The total flux was approximately  $(1.51 \pm 0.03) \times 10^{-7} \text{ cm}^{-2} \text{s}^{-1}$  for  $E_n > 1 \text{ MeV}$ without the polyethylene shielding. The results for the residual neutron-induced background with the copper and PE shielding setups are shown in Fig. 7. We observed that the environmental neutrons contribute approximately  $1.6 \times 10^{-3}$ cts/keV/kg/yr in the <sup>100</sup>Mo  $0\nu\beta\beta$  ROI with "*multiplicity* = 1" cut applied. It should be noted that there are currently no published experimental data on CJPL environmental neutron flux exceeding 20 MeV. Thus, high-energy neutrons ( $E_n >$ 20 MeV) were not considered in our simulations. Nevertheless, according to existing simulation studies performed for LNGS, the integral flux exceeding 10 MeV (mainly from muon-induced processes) is three orders of magnitude lower than that below 10 MeV (mainly from radioactivity-induced processes) [38, 41]. For the CJPL, owing to the larger depth, the flux of cosmic muon-induced neutrons was even lower. The relative background contribution from  $E_{\rm n} > 20 \text{ MeV}$ high-energy neutron was expected to be low.

# 3.4 Underground cosmic muon-induced background

Cosmogenic backgrounds play a pivotal role in rare-event search experiments [39]. At Earth's surface, cosmic rays



Fig. 7 (Color online) Simulated results of the neutron-induced background

predominantly consist of high-energy muons [40]. The significant rock overburden of CJPL (equivalent to 6700 m of water) dramatically reduces the cosmic muon flux underground-by about eight orders of magnitude compared to the flux at sea level. As a result, the cosmogenic background does not pose a significant threat to the Jinping bolometric experiments. Nevertheless, for the heightened sensitivity objectives of upcoming  $0\nu\beta\beta$  experiments, a thorough evaluation of the residual muon-induced background is imperative.

To estimate the muon-induced background level, it is essential to understand both the energy and angular distributions of underground muons in a given experimental cavern. The muon energy was derived using a parametric function [41]:

$$\frac{\mathrm{d}N}{\mathrm{d}E_{\mu}} = Ce^{-bh(\gamma_{\mu}-1)} \left( E_{\mu} + \varepsilon_{\mu} \left( 1 - e^{-bh} \right) \right)^{-\gamma_{\mu}},\tag{2}$$

where b = 0.4/km·w·e,  $\gamma_{\mu} = 3.77$ , and  $E_{\mu} = 693$  GeV were used, as suggested.

Figure 8 shows the underground muon spectrum from a Geant4-based simulation study and compares it with the Mei–Hime parameterization results [42]. The parametric function accurately described the Geant4 simulation results. For the angular distribution, additional information regarding rock overburden is required to determine the flux in a specific direction. In this study, the zenith and azimuthal angle distributions of the underground muons were sampled according to JNE measurements [43]. The experimental distributions can be described using the parametric function discussed in Ref. [44]. The total underground



**Fig.8** (Color online) Energy spectrum of the CJPL muons. Results from the parametric function are shown in comparison with that from the Monte Carlo simulation

muon flux used as the input for the simulation was  $3.53 \times 10^{-10}$  cm<sup>-2</sup>s<sup>-1</sup> based on the measurement at CJPL Phase II.

The simulated results of the muon-induced background contributions are shown in Fig. 9. We observed that the background contribution of muon events, including muon-induced secondaries, corresponds to a low level of  $2.2 \times 10^{-5}$  cts/keV/kg/yr in the <sup>100</sup>Mo  $0\nu\beta\beta$  ROI. It was also observed that the multiplicity anti-coincidence cut significantly suppressed the muon-induced background by an order of magnitude.

## 4 Conclusion

We conducted a simulation study evaluating the background shielding for the Jinping 100 Mo-based bolometric demonstration experiment aimed at searching for neutrinoless doublebeta decay. Having established the shielding arrangement through an attenuation study, we assessed the environmental gamma, neutron, and muon-induced backgrounds. Our findings indicate that within the <sup>100</sup>Mo  $0\nu\beta\beta$  ROI, gamma rays and neutrons contribute to background levels of approximately  $1.3 \times 10^{-3}$  cts/keV/kg/yr and  $1.6 \times 10^{-3}$  cts/keV/kg/yr, respectively. The underground cosmic ray muon's contribution was minor, only reaching  $2.2 \times 10^{-5}$  cts/keV/kg/yr in the ROI when the anti-coincidence cut was applied. Additionally, given the bolometer's slow data acquisition rate, potential pile-up events, notably coincident low-energy gamma rays, could also add to the discernible experimental background. It is worth noting that our simulations were done for a demonstration experiment without supplementary passive background shielding. In CJPL Phase II, a meticulously designed combined shielding apparatus (comprising 20 cm of lead and 100 cm of PE) was implemented for radioactive



Fig. 9 (Color online) Simulated results of the muon-induced background contribution

background protection. We anticipate a significant reduction in the combined contributions from environmental gamma and neutron backgrounds to levels below  $10^{-3}$  cts/keV/kg/y, aligning with our objectives for this demonstration experiment. For the  $0\nu\beta\beta$  experiment, background plays a pivotal role, directly influencing experimental sensitivity. Therefore, curbing background contribution emerges as a critical facet of bolometric experimental studies and advancements. Regarding the environmental background, precise in situ measurements of both neutron and gamma spectra, extending into high-energy regions, are paramount for designing shielding systems tailored to future high-sensitivity  $0\nu\beta\beta$ research endeavors at CJPL.

Author Contributions All authors contributed to the study conception and design. Material preparation, data collection, and analysis were performed by Wei Chen, Long Ma, Jin-Hui Chen, and Huan-Zhong Huang. The first draft of the manuscript was written by Long Ma, and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

**Data Availability** The data that support the findings of this study are openly available in Science Data Bank at https://doi.org/10.57760/scien cedb.10675 and https://cstr.cn/31253.11.sciencedb.10675.

## Declarations

**Conflict of interest** Jin-Hui Chen is an editorial board member for Nuclear Science and Techniques and was not involved in the editorial review, or the decision to publish this article. All authors declare that there are no competing interests.

# References

- S.M. Bilenky, C. Giunti, Neutrinoless double-beta decay: a probe of physics beyond the Standard Model. Int. J. Mod. Phys. A 30, 1530001 (2015). https://doi.org/10.1142/S0217751X1530001X
- D.Q. Adams, C. Alduino, K. Alfonso et al., (CUORE collaboration), improved limit on neutrinoless double-beta decay in <sup>130</sup>Te with CUORE. Phys. Rev. Lett. **124**, 122501 (2020). https://doi.org/10.1103/PhysRevLett.124.122501
- J.B. Albert, G. Anton, I. Badhrees et al., (EXO collaboration), search for neutrinoless double-beta decay with the upgraded EXO-200 detector. Phys. Rev. Lett. **120**, 072701 (2018). https://doi.org/ 10.1103/PhysRevLett.120.072701
- M. Agostini, A.M. Bakalyarov, M. Balata et al., (GERDA collaboration), improved limit on neutrinoless double-*β* decay of <sup>76</sup>Ge from GERDA phase II. Phys. Rev. Lett. **120**, 132503 (2018). https://doi.org/10.1103/PhysRevLett.120.132503
- A. Gando, Y. Gando, T. Hachiya et al., (KamLAND-Zen collaboration), Search for Majorana neutrinos near the inverted mass hierarchy region with KamLAND-Zen. Phys. Rev. Lett. 117, 082503 (2016). https://doi.org/10.1103/PhysRevLett.117.082503
- M. Biassoni, O. Cremonesi, Search for neutrino-less double beta decay with thermal detectors. Prog. Part. Nucl. Phys. **114**, 103803 (2020). https://doi.org/10.1016/j.ppnp.2020.103803
- W.R. Armstrong, et al. (The CUPID Interest Group), arXiv:1907. 09376
- E. Armengaud, C. Augier, A.S. Barabash et al., (CUPID collaboration), New limit for neutrinoless double-beta decay of <sup>100</sup>Mo

from the CUPID-Mo experiment. Phys. Rev. Lett. **126**, 181802 (2020). https://doi.org/10.1103/PhysRevLett.126.181802

- E. Armengaud, C. Augier, A. Barabash et al., Precise measurement of 2νββ decay of <sup>100</sup>Mo with the CUPID-Mo detection technology. Eur. Phys. J. C 80, 674 (2020). https://doi.org/10.1140/epjc/s10052-020-8203-4
- A. Alessandrello, C. Arpesella, C. Brofferio et al., Measurements of internal radioactive contamination in samples of Roman lead to be used in experiments on rare events. Nucl. Instrum. Meth. Phys. Res. B 142, 163–172 (1998). https://doi.org/10.1016/S0168-583X(98)00279-1
- A. Luqman, D.H. Ha, J.J. Lee et al., Simulations of background sources in AMORE-I experiment. Nucl. Instrum. Meth. Phys. Res. A 855, 140–147 (2017). https://doi.org/10.1016/j.nima.2017.01. 070
- F. Alessandria, R. Ardito, D.R. Artusa et al., (CUORE collaboration), Validation of techniques to mitigate copper surface contamination in CUORE. Astropart. Phys. 45, 13–22 (2013). https://doi. org/10.1016/j.astropartphys.2013.02.005
- H.W. Bae, E.J. Jeon, Y.D. Kim et al., Neutron and muon-induced background studies for the AMoRE double-beta decay experiment. Astropart. Phys. **114**, 60–67 (2020). https://doi.org/10. 1016/j.astropartphys.2019.06.006
- C. Alduino, K. Alfonso, D.R. Artusa et al., (CUORE collaboration), The projected background for the CUORE experiment. Eur. Phys. J. C 77, 543 (2017). https://doi.org/10.1140/epjc/ s10052-017-5080-6
- W. Chen, L. Ma, J.H. Chen et al., Cosmogenic background study for a <sup>100</sup>Mo-based bolometric demonstration experiment at China JinPing underground Laboratory. Eur. Phys. J. C 82, 549 (2022). https://doi.org/10.1140/epjc/s10052-022-10501-y
- P.N. Peplowski, J.T. Wilson et al., Cosmogenic radionuclide production modeling with Geant4: experimental benchmarking and application to nuclear spectroscopy of asteroid (16) Psyche. Nucl. Instrum. Meth. Phys. Res. B 446, 43–57 (2019). https://doi.org/ 10.1016/j.nimb.2019.03.023
- F. Bellini, C. Bucci, S. Capelli et al., Monte Carlo evaluation of the external gamma, neutron and muon induced background sources in the CUORE experiment. Astroparticle Physics 33, 169–174 (2010). https://doi.org/10.1016/j.astropartphys.2010. 01.004
- M. Haffke, L. Baudis, T. Bruch et al., Background measurements in the Gran Sasso underground laboratory. Nucl. Instrum. Meth. Phys. Res. A 643, 36–41 (2011). https://doi.org/10.1016/j.nima. 2011.04.027
- J. Allison, K. Amako, J. Apostolakis et al., Recent developments in Geant4. Nucl. Instrum. Methods A 835, 186 (2016). https://doi. org/10.1016/j.nima.2016.06.125
- S. Agostinelli, J. Allison, K. Amako et al., Geant4—a simulation toolkit. Nucl. Instrum. Methods A 506, 250 (2003). https://doi. org/10.1016/S0168-9002(03)01368-8
- D.-J. Zhao, S. Feng, P.-J. Cheng et al., Conceptual design of a Cs<sub>2</sub>LiLaBr<sub>6</sub> scintillator-based neutron total cross section spectrometer on the back-n beam line at CSNS. Nucl. Sci. Tech. 34, 3 (2023). https://doi.org/10.1007/s41365-022-01152-5
- L.-H. Zhou, S.-Y. Cao, T. Sun et al., A refined Monte Carlo code for low-energy electron emission from gold material irradiated with sub-keV electrons. Nucl. Sci. Tech. 34(4), 54 (2023). https:// doi.org/10.1007/s41365-023-01204-4
- Q.D. Hu, H. Ma, Z. Zeng et al., Neutron background measurements at China Jinping underground laboratory with a Bonner multi-sphere spectrometer. Nucl. Instrum. Meth. Phys. Res. A 859, 37–40 (2017). https://doi.org/10.1016/j.nima.2017.03.048
- Y. Elmahroug, B. Tellili, C. Souga, Calculation of gamma and neutron shielding parameters for some materials polyethylenebased. Int. J. Phys. Res. 3(1), 33–40 (2013)

- 25. D. Venkata Subramanian, A. Haridas, D. Sunil Kumar, Neutron attenuation studies with borated polyethylene slabs containing 30% natural boron and its comparison with hydrogenous materials. Indian J. Pure Appl. Phys 56, 583–586 (2018)
- H. Ma, Z. She, W.H. Zeng et al., In-situ gamma-ray background measurements for next generation CDEX experiment in the China Jinping underground laboratory. Astropart. Phys. **128**, 102560 (2021). https://doi.org/10.1016/j.astropartphys.2021.102560
- Z. Zeng, J. Su, H. Ma et al., Environmental gamma background measurements in China Jinping Underground Laboratory. J. Radioanal. Nucl. Chem. **301**(2), 443–450 (2014). https://doi.org/10. 1007/s10967-014-3114-1
- Y.P. Shen, J. Su, W.P. Liu et al., Measurement of γ detector backgrounds in the energy range of 3–8 MeV at Jinping underground laboratory for nuclear astrophysics. Sci. China Phys. Mech. Astron. 60(10), 102022 (2017). https://doi.org/10.1007/ s11433-017-9049-3
- Q. Yue, C.J. Tang, J.P. Cheng et al., A Monte Carlo study for the shielding of γ backgrounds induced by radionuclides for CDEX. Chin. Phys. C 35, 282 (2011). https://doi.org/10.1088/1674-1137/ 35/3/013
- R.M.J. Li, S.K. Liu, S.T. Lin et al., Identification of anomalous fast bulk events in a p-type point-contact germanium detector. Nucl. Sci. Tech. 33(5), 57 (2022). https://doi.org/10.1007/ s41365-022-01041-x
- B. Schmidt et al., CUPID-Mo Collaboration, First data from the CUPID-Mo neutrinoless double beta decay experiment. J. Phys. Conf. Ser. 1468(1), 012129 (2020). https://doi.org/10.1088/1742-6596/1468/1/012129
- B. Aharmim, S.N. Ahmed, A.E. Anthony et al., Cosmogenic neutron production at the sudbury neutrino observatory. Phys. Rev. D 100, 112005 (2019). https://doi.org/10.1103/PhysRevD.100. 112005
- V. A. Kudryavtsev, N.J.C. Spooner, J. E. McMillan, Simulations of muon-induced neutron flux at large depths underground. arXiv: hep-ex/0303007v1 (2003)
- Z.M. Zeng, H. Gong, Q. Yue et al., Thermal neutron background measurement in CJPL. Nucl. Instrum. Meth. A 804, 108–112 (2015). https://doi.org/10.1016/j.nima.2015.09.043

- G. Bruno, W. Fulgione, Flux measurement of fast neutrons in the Gran Sasso underground laboratory. Eur. Phys. J. C 79, 747 (2019). https://doi.org/10.1140/epjc/s10052-019-7247-9
- Q.H. Wang, A. Abdukerim, W. Chen et al., An improved evaluation of the neutron background in the PandaX-II experiment. Sci. China Phys. Mech. Astron. 63(3), 231011 (2020). https://doi.org/ 10.1007/s11433-019-9603-9
- Q. Du, S.T. Lin, S.K. Liu et al., Measurement of the fast neutron background at the China Jinping underground laboratory. Nucl. Instrum. Meth. Phys. Res. A 889, 105–112 (2018). https://doi.org/ 10.1016/j.nima.2018.01.098
- H. Wulandari, J. Jochum, W. Rau et al., Neutron flux at the Gran Sasso underground laboratory revisited. Astropart. Phys. 22, 313– 322 (2004). https://doi.org/10.1016/j.astropartphys.2004.07.005
- E. Andreotti, C. Arnaboldi, F.T. Avignone III et al., Muon-induced backgrounds in the CUORICINO experiment. Astropart. Phys. 34, 18–24 (2010). https://doi.org/10.1016/j.astropartphys.2010.04.004
- T.K. Gaisser, R. Engel, E. Resconi, Cosmic rays and particle physics. (Cambridge University Press, New York, 1990), p. 71
- D.M. Mei, A. Hime, Muon-induced background study for underground laboratories. Phys. Rev. D 73, 053004 (2006). https://doi. org/10.1103/PhysRevD.73.053004
- 42. X.H. Hu, Simulation of cosmic ray background at CJPL, Master's Dissertation, Nankai University, (2013) (in Chinese)
- Z.Y. Guo, L.B. Peters, S.M. Chen et al., Muon flux measurement at China Jinping underground laboratory. Chin. Phys. C 45, 025001 (2021). https://doi.org/10.1088/1674-1137/abccae
- H. Arslan, M. Bektasoglu, Geant4 simulation study of deep underground muons: Vertical intensity and angular distribution. Adv. High Energy Phys. 2013, 391573 (2013). https://doi.org/10.1155/ 2013/391573

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.